

























Named lava flows:	Copper Harbor Conglomerate	Unnamed rock sequences:
		
		Numerous thin ophitic flows with as many as seven interbedded sandstone and conglomerate units (see fig. 17 for details)
Scoville Point Flow (porphyrite) (psp)		Several thin ophitic flows
Edwards Island Flow (trap) (pei)		
Middle Point Flow (porphyrite) (pmp)		Conglomerate—known from drill records
		Several thin ophitic flows
Long Island Flow (trap) (pli)		
		Sandstone—known from drill records
Tobin Harbor Flow (porphyrite) (pth)		
Washington Island Flow (ophite) (pwi)		Tuff-breccia
		Tuff-breccia
Greenstone Flow (ophite) (pg)		
		Conglomerate—known from drill records
Grace Island Flow (porphyrite) (pgi)		
		Sequence of thin to thick (more than 100 ft) flows chiefly ophitic, with one or more sedimentary units suggested by drill data
		Tuff-breccia
Minong Flow (trap) (pm)		
		Sequence of thin to thick flows, chiefly ophitic, with one or more sedimentary units suggested by drill data
Huginnin Flow (porphyrite) (ph)		
		One or more ophitic flows present locally
Hill Point Flow (ophite) (php)		
		Sequence of thin to moderately thick flows, chiefly ophitic. Several sedimentary units and a felsite indicated by drill records
		Breccia
Amygdaloid Island Flow (trap) (pai)		Lava flows, chiefly ophitic