



# Hemlock Woolly Adelgid



Eastern hemlock trees are some of the largest and most common trees in the Great Smoky Mountains. Unfortunately, they are under attack from a non-native insect called the hemlock woolly adelgid. Without successful intervention, the hemlock woolly adelgid is likely to kill most of the hemlock trees in the park.

Called the "redwood of the east," eastern hemlocks (*Tsuga canadensis*) can grow more than 150 feet tall on trunks measuring six feet in diameter. Some hemlocks in the park are over 500 years old.

Over 800 acres of old-growth hemlock trees grow in the Smokies—more than in any other national park. Younger hemlock forests cover an additional 90,000 acres of land in the park. Originally discovered here in 2002, adelgid infestations have now spread throughout the park's hemlock forests. In some areas infested trees have already begun to die.

## WHAT TO LOOK FOR

The hemlock woolly adelgid (*Adelges tsugae*) is an aphid-like insect that covers itself with a white, waxy "wool" which acts as a protective coating for the insect. Adelgid infestations are easily recognizable by the appearance of tiny "cotton balls" at the base of hemlock needles. The "wool" is most conspicuous on the undersides of branches from fall through spring.

The hemlock woolly adelgid feeds on the sap at the base of hemlock needles, disrupting nutrient flow and causing the needles to change from deep green to a grayish green, then fall off. Without needles the tree starves to death, usually within three to five years of the initial attack.

## WHAT IS THE NATIONAL PARK DOING?

Three different types of treatments are used to kill adelgids on hemlock trees:

### Foliar Treatments

Hemlocks in developed areas and backcountry sites accessible by road are treated with insecticidal soap or horticultural oils. Sprayed from truck-mounted spray units, these sprays smother and dry-out the adelgids on contact. The equipment can spray up to 80 feet into the canopy of large roadside trees and can efficiently treat areas of smaller trees. However this method kills only the insects that are present on the tree at the time of application and requires retreatment every six months to one year.

### Systemic Treatments

Hemlocks that are growing near campsites or are too tall to be sprayed are treated with a systemic insecticide (imidacloprid) either by soil drenching or by injecting the insecticide directly into the trunk. Soil drenching involves temporarily removing the duff—a layer of organic matter—from around the base of the tree then pouring a mixture of imidacloprid and water on the bare ground around the tree within a foot of the trunk. The duff layer is then replaced. The results of insecticidal treatments have been dramatic. Trees with ashen gray foliage prior to treatment recover their color and produce new growth. Treatments may remain effective for up to three years.

### Predator Beetles

The park began releasing predatory beetles, which feed exclusively on adelgids, as a biocontrol in 2002. It will take several years before populations of beetles increase enough to control adelgid infestations. Although it is too early to assess the overall success of this biocontrol, preliminary monitoring results are encouraging.

## YOU CAN HELP

Although the adelgid will forever alter hemlock forests of the Smokies, with continued funding, dedicated staff, and committed partners, future visitors to the park will still be able to marvel at the "redwood of the east."

Efforts to control hemlock woolly adelgids are being funded through the Save the Hemlocks initiative of the Friends of the Smokies, a nonprofit organization. For additional information, call (865) 453-2428 or visit [www.friendsofthesmokies.org](http://www.friendsofthesmokies.org)

## LEARN MORE ABOUT THE ADELGID

Learn more about the hemlock woolly adelgid by visiting [www.saveourhemlocks.org](http://www.saveourhemlocks.org) or call (865) 436-1252.