





**Common Native Plants**



**Wild Ginger**  
*Asarum caudatum*

Distinct heart-shaped leaves form small carpets in moist shady areas.  
 Aromatic stems and roots.



**Redwood Sorrel**  
*Oxalis oregana*

Carpets forest floor.  
 Flowers white to pinkish.  
 Leaves in 3 heart shaped sections.




**Salal**  
*Gaultheria shallon*

Waxy green leaves with sticky stems.  
 Flowers hanging white bells.  
 Berries dark purple.




**Thimbleberry**  
*Rubus parviflorus*

Tall shrub. Leaves 5-lobed, hairy and toothed.  
 Fruit red, thimble shaped.  
 Ripe early-mid summer.




**Poison Oak**  
*Toxicodendron diversilobum*

Grows as shrub or vine.  
 Leaves grow in groups of three. Green in summer turning red in fall. **CAUTION:** Do not touch, can cause allergic reaction, skin rash.




**Salmonberry**  
*Rubus spectabilis*

Common Shrub along creeks and edges of forests. Leaves serrated. Stalk has prickles.  
 Produces yellow to red berries.




**Red Huckleberry**  
*Vaccinium parvifolium*

Deciduous, tall shrub with soft green leaves in summer.  
 Bark bright green.  
 Berries red, small, ripe in mid-summer.



**Evergreen Huckleberry**  
*Vaccinium ovatum*

Tall evergreen shrub with waxy, toothed, green leaves.  
 Berries black, clustered, ripe in fall.




**Trail Plant**  
*Adenocaulon bicolor*

Leaves smooth and green on top  
 Surface, fuzzy and white on underside. When trampled, the white arrow-shaped undersides point in direction of the disturber.




**Common Native Trees**



**Tanoak**

*Lithocarpus densiflorus* var. *densiflorus*


Leaves leathery, toothed.  
 Bark grayish brown.  
 Acorns hairy-capped and are important  
 Source of food for wildlife.



**Red Alder**

*Alnus rubra*


Grows in open and damp habitats.  
 Leaves coarsely toothed.  
 Bark white-gray.



**California Laurel**

(Pepperwood, Bay)  
*Umbellularia californica*

Leaves shiny green, yellow in  
 fall, aromatic when rubbed in  
 hands.



**Douglas-fir**

*Pseudotsuga menziesii* var. *menziesii*


Found on ridges.  
 Bark brown and furrowed, brittle.  
 Often carries more  
 lichen than Coast Redwoods.



**Big-leaf Maple**

*Acer macrophyllum*


Grows along streams.  
 Bark covered with mosses.  
 Leaves yellow in fall.



**Sitka Spruce**

*Picea sitchensis*

Grows close to coast.  
 Needles grow on all sides of twig.  
 Bark gray, chip-like.



**Coast Redwood**

*Sequoia sempervirens*  
 (sun needles)

Fibrous reddish-brown bark, deep  
 furrows extend up tree. Cones  
 olive sized.



**Coast Redwood**

*Sequoia sempervirens*  
 (shade needles)

Scaly needles (left) from top of tree.  
 Flat needles (shown) from lower  
 reaches where light is  
 less prevalent.



**Western Hemlock**

(Octopus Tree)  
*Tsuga heterophylla*

Often grows on  
 decaying logs and stumps.  
 Drooping branches with soft  
 needles.