


Geologic Overview of a Bison-Carcass Site at Norris Geyser Basin, March 2004



Cover Photograph. Overview of the bison carcasses along the Gibbon River.
by

Hank Heasler and Cheryl Jaworowski
Geology Program
Yellowstone Center for Resources
Yellowstone National Park

With assistance from
Kerry Gunther, Susan Chin, Travis Wyman
Bear Management Program
Yellowstone Center for Resources
Yellowstone National Park

March 23, 2004

Introduction

On March 10, 2004, Bear Management personnel noticed an unusual grouping and position of bison carcasses near Norris Geyser Basin. On March 11, 2003, YNP geologists joined bear management biologists (Travis Wyman and Susan Chin) to investigate 5 bison along the Gibbon River. The adults and calves were lying on their sides with their feet perpendicular to their bodies (Figure 1). The unusual position of the carcasses led biologists to suspect that the bison had died very rapidly, as a group.



Biologists established that the bison had been dead for approximately one week. Reconnaissance of the area showed that the 5 bison died along the Gibbon River and downstream of multiple gas vents. Areas with multiple gas vents are associated with thermally baked ground, minimal vegetation and sulfur deposits. Multiple gas vents occur uphill from the bison carcasses on both sides of the Gibbon River.

Figure 1. Bison lying on its side with feet perpendicular to its body. All 5 bison were lying in a similar position.

Data

Bison Management personnel (Doug Blanton and others) visited the site on March 10th, collecting incisors and tail hair. Members of the Bear Management team examined the 5 bison on March 11th and collected samples of blood, lung, liner and windpipe. The Bear Management team also examined body fat and bone marrow for each carcass. Body fat was present on all bison. Bone marrow was pink and solid for 4 bison. One calf had red and gelatinous bone marrow.

Table 1. Bison

Number	Animal	Age	Tissue	Teeth	Body Fat
1	Bison	Calf	No	Yes	Yes
2	Bison	Adult	No	Yes	Yes
3	Bison	Adult	No	Yes	Yes
4	Bison	Yearling	No	Yes	Yes
5	Bison	Calf	Yes	Yes	Yes

Within the thermal area, an Orion Multi-Gas Detector measured concentrations of hydrogen sulfide and oxygen. Concentrations of hydrogen sulfide (H₂S) varied from a high of greater than 200 ppm to 0 ppm (Figures 2 and 3, Table 2). Concentrations of greater than 200 ppm H₂S were found at a vent immediately upstream from the bison carcasses (Figures 2 and 3). A second vent in the thermal area emitted steam with 177 ppm of H₂S (Table 2). Within the thermal area, the air contained 56 ppm of H₂S (Table 2). On the top of a small hill (Table 2), the gas meter measured 0 ppm H₂S in the gentle wind. Downstream from the thermal area, the gas meter measured 1 ppm H₂S (Table 2). Oxygen varied from 16 % (within a vent) to 20.8 % (within the air). On the morning of March 11th, there was a gentle breeze mixing the air.




Figure 2. Photograph of the thermal area upstream from the bison carcasses. Brown, yellow and white colors indicate thermally baked ground and deposition of minerals. A H₂S concentration of 56 ppm (Table 2) was measured in the air of this thermal area on a breezy March morning.



Figure 3. Close-up of steam/gas vent in Figure 2 and gas meter. H₂S emissions from this vent were greater than 200 ppm. Notice the yellow sulfur crystals around the vent and the red-brown, thermally baked ground. The vent shown in this photograph is in Table 2.

An Oakton temperature meter with a Type K thermocouple measured ground and water temperatures. Within the brown, tan, and yellow ground, temperatures varied from 27°C to 93.4°C at 4-cm depth. In places, yellow crystals of native sulfur were visible around vents. Water temperatures of a small seep (13.5°C) and thermal spring (55.9°C) were greater than the temperature of the Gibbon River (7.9°C) in the area of the bison carcasses.

Table 2. Gas Concentrations

Number	Feature	H ₂ S (ppm)	O ₂ (%)	Temperature (°C)
1	Gibbon River	Not Meas.	Not Meas.	7.9
2	Thermal Spring	Not Meas.	Not Meas.	55.9
3	Thermal Pools	1	Not Meas.	13.5
4	Mud Pot	177	20.6	82.5
5	Ground	Not Meas.	Not Meas.	93.4
6	Gas Vent	>200	16.0	Not Meas.
7	Gas Vent	>200	Not Meas.	83.0
8	Air	56	20.8	Not Meas.
9	Hill Top	0	20.8	Not Meas.

Discussion

In the field, the YCR team of geologists and biologists discussed the possible effects of toxic gases such as H₂S. Hydrogen sulfide (H₂S) is easily recognized at concentrations of 1 ppm. The odor of H₂S increases up to 30 ppm. From 1-30 ppm, humans recognize its characteristic rotten-egg smell. From 30-100 ppm, humans recognize H₂S gas as “sickeningly sweet”. Above 100 ppm, a human’s ability to detect H₂S is impaired. Prolonged exposures can lead to bronchitis, pneumonia, headaches, irritations of eyes, nose or throat, dizziness, nausea, shock, convulsions, coma, or death.

H₂S is classed as a *chemical asphyxiant*, similar to carbon monoxide and cyanide gases. It inhibits cellular respiration and uptake of oxygen, causing biochemical suffocation. With a vapor density of 1.19, hydrogen sulfide is approximately 20 percent heavier than air, so this invisible gas may collect in depressions in the ground and in confined spaces.

Another toxic gas that occurs in geothermal areas is carbon dioxide (CO₂). At 10%, carbon dioxide paralyzes a human’s respiratory system. The result is fatal with no evidence of an agonal struggle. The ratio of carbon dioxide to hydrogen sulfide in most geothermal areas ranges between 1,000 to 1 to 100 to 1 (Bill Evans, United States Geological Survey, Personal Communication, March, 2004).

Hydrogen sulfide and carbon dioxide can accumulate in topographically low areas on cold, calm nights because they are denser than air. For Norris Geyser Basin, no weather station exists. However, an electronic temperature logger deployed at the Norris Museum shows unusually cold temperatures on the evening of March 1st and early morning of March 2nd, 2004 (Figure 4, -17°C, (1°F) at 2:56 a.m. MST). Also, the elevation of the mapped gas vent is approximately 20 to 30 feet higher in elevation than the elevation of

the bison carcasses. Therefore, it is possible that the 5 bison were asphyxiated by CO₂ and/or H₂S gases on a cold night with still air in early March.




Figure 4. Graph of air temperature at Norris Museum from February 25th to March 11th, 2004.

The death of animals by toxic gases is not unusual in Yellowstone National Park. Weed (1889), noted the death of 6 bears, one elk, and “several squirrels, rock hares, and other small mammals, besides numerous dead butterflies and insects” in an area now called Death Gulch in the upper Lamar River valley. Jagger (1899) also wrote of natural toxic gases in the northeast corner of Yellowstone and noted 7 dead bears in Dead Gulch in the late 1800’s. Ravines or topographically low areas can accumulate the dense, toxic gases during times of still air. A cursory survey of holdings at the Yellowstone Research Library indicates that many other people have recognized the dangers of toxic gases within Yellowstone (Ash, 1939; Frisbee, 1960, Fournier, 1959; Love and Good, 1970; Traphagen, 1904).

Conclusions

Yellowstone National Park’s active geothermal system produces toxic gases such as hydrogen sulfide (H₂S) and carbon dioxide (CO₂). Because few of Yellowstone’s thermal areas are in closed depressions that would concentrate these gases, the likelihood of a fatal exposure is remote. Historically, people have observed animals killed by natural, toxic emissions of gases.

Gas measurements and field reconnaissance support the idea that bison were overwhelmed by a toxic emission of gas. Hydrogen sulfide (H₂S) and/or carbon dioxide (CO₂) are considered the most likely gases to have catastrophically affected the 5 bison near the Gibbon River. Being more dense than air, hydrogen sulfide and /or carbon dioxide gases from vents in the nearby thermal area could have drained downhill towards the Gibbon River and overwhelmed 5 bison during late evening of March 1 or early morning, March 2, 2004.

Recommendations

A reconnaissance of gases will be undertaken in the entire Norris area. This will include random samples of hydrogen sulfide, carbon dioxide, and oxygen. Both air and vent samples will be taken. All vent samples will include temperature measurements and a digital photograph.

YCR Geology personnel have been unable to locate any Yellowstone National Park carbon dioxide measuring equipment. Consequently, a hand-held CO₂ measuring unit will be purchased to complete a reconnaissance of CO₂ gas.

References Cited

Ash, S.H., 1939, Investigation of gas hazards at proposed bridge foundation sites in Yellowstone National Park: Yellowstone National Park, Yellowstone Research Library, vertical file, Book, 48 p.

Frisbee, R., 1960, Report on Death Gulch: Yellowstone National Park, Yellowstone Research Library, vertical file, 3 p.

Fournier, R.O., 1959, Death Gulch Gas Analysis: Yellowstone National Park, Yellowstone Research Library, vertical file, Book.

Hydrogen Sulfide Safety Factsheet. http://www.ppm-technology.com/factsheets/h2s_safety.html

Jaggar, T.A., 1899, Death Gulch-A natural bear-trap: Appleton's popular science monthly, vol. 54, no. 4, p.475-481.

Love, J.D., and Good, J.M., 1970, Hydrocarbons in thermal areas, Northwestern Wyoming: U.S. Geological Survey Professional Paper 644-B, 23 p.

Traphagen, F.W., 1904, Death Gulch: Science, vol. 19, no. 485, p. 632-634.

Weed, W.H., 1889, A deadly gas-spring in the Yellowstone Park: Science vol. 13, no. 315, p.130-132.