

TABLE 13. Volumetric percentages of forage in rumens of 25 white-tailed deer collected from June 1972 to April 1974 in Big Bend National Park.^a

Species	Summer (3)	Late summer (6)	Winter (9)	Spring (7)	Total years (25)
Browse species					
<i>Acacia</i> spp.	2.7	12.2	7.4	5.1	7.4
<i>Quercus</i> spp.	1.3	11.0	4.2	8.4	6.7
<i>Rhus virens</i>	—	0.7	12.1	5.0	5.9
<i>Porlieria angustifolia</i>	0.3	12.5	1.6	—	3.6
<i>Garrya wrightii</i>	—	10.5	—	—	2.5
<i>Rhus microphylla</i>	3.7	—	—	5.1	1.9
<i>Prosopis glandulosa</i>	—	5.8	—	—	1.4
<i>Dalea</i> spp.	2.3	3.2	0.6	—	1.2
<i>Rhus</i> spp.	—	0.5	2.3	—	1.0
<i>Cercocarpus</i> spp.	3.7	1.2	0.6	—	0.9
<i>Rhus trilobata</i>	—	1.5	0.7	—	0.6
<i>Dyssodia</i> spp.	—	0.3	0.9	T ^b	0.4
<i>Fraxinus greggii</i>	—	—	—	1.4	0.4
<i>Nolina erumpens</i>	1.7	—	—	—	0.2
<i>Fallugia paradoxa</i>	—	—	—	0.9	0.2
<i>Diospyros texana</i>	1.3	—	0.4	—	0.3
<i>Forestiera angustifolia</i>	—	—	—	0.4	0.1
<i>Phoradendron</i> spp.	—	—	—	0.1	T
<i>Viguiera</i> spp.	1.0	—	—	0.3	0.2
<i>Juniperus</i> spp.	0.3	—	0.6	—	0.2
<i>Pinus cembroides</i>	—	0.2	—	—	T
<i>Xanthocephalum</i> spp.	—	—	—	—	—
<i>Prunus havardii</i>	—	—	—	—	—
<i>Hoffmanseggia</i> spp.	—	—	—	—	—
<i>Leucophyllum</i> spp.	—	—	—	—	—
Total browse	18.3	59.6	31.4	26.7	35.1
Forb species					
<i>Euphorbia serrula</i>	18.7	3.7	4.4	6.6	6.6
<i>Erigeron divergens</i>	—	1.3	5.4	—	2.3
<i>Lotus oroboides</i>	—	—	5.1	—	1.8
<i>Eriogonum</i> spp.	—	—	—	5.3	1.5
<i>Verbena</i> spp.	1.0	2.5	0.6	—	0.9
<i>Argythamnia humilis</i>	—	—	1.7	—	0.6
<i>Menodora</i> spp.	0.7	—	0.4	—	0.2
<i>Penstemon havardii</i>	—	—	—	0.4	0.1
<i>Hedyotis</i> spp.	—	—	0.2	—	0.1
<i>Artemisia</i> spp.	—	0.2	0.1	—	0.1
<i>Setcreasea</i> spp.	—	0.3	—	—	0.1
<i>Mentha</i> spp.	0.3	—	—	—	T
<i>Croton</i> spp.	—	—	—	T	T
Total forbs	20.7	8.0	17.9	12.3	14.3
Succulent species					
<i>Agave lecheguilla</i>	8.0	3.2	20.9	27.6	17.0