



OREGON GEOLOGY

published by the
Oregon Department of Geology and Mineral Industries



VOLUME 58, NUMBER 3

MAY 1996



IN THIS ISSUE:

Reconstructions of Eocene and Oligocene Plants and Animals of Central Oregon

OREGON GEOLOGY

(ISSN 0164-3304)

VOLUME 58, NUMBER 3

MAY 1996

Published bimonthly in January, March, May, July, September, and November by the Oregon Department of Geology and Mineral Industries. (Volumes 1 through 40 were entitled *The Ore Bin*.)

Governing Board

John W. Stephens, Chair..... Portland
Jacqueline G. Haggerty..... Weston Mountain
Ronald K. Culbertson..... Myrtle Creek

State Geologist..... Donald A. Hull
Deputy State Geologist..... John D. Beaulieu
Publications Manager/Editor..... Beverly F. Vogt
Production Editor..... Klaus K.E. Neuendorf
Production Assistants..... Geneva Beck
Kate Halstead

Main Office: Suite 965, 800 NE Oregon Street # 28, Portland 97232, phone (503) 731-4100, FAX (503) 731-4066.

Baker City Field Office: 1831 First Street, Baker City 97814, phone (541) 523-3133, FAX (541) 523-5992.

Mark L. Ferns, Regional Geologist.

Grants Pass Field Office: 5375 Monument Drive, Grants Pass 97526, phone (541) 476-2496, FAX (541) 474-3158.

Thomas J. Wiley, Regional Geologist.

Mined Land Reclamation Program: 1536 Queen Ave. SE, Albany 97321, phone (541) 967-2039, FAX (541) 967-2075.

Gary W. Lynch, Supervisor.

The Nature of the Northwest Information Center: Suite 177, 800 NE Oregon Street # 5, Portland, OR 97232-2162, phone (503) 872-2750, FAX (503) 731-4066, Donald J. Haines, Manager.

Second class postage paid at Portland, Oregon. Subscription rates: 1 year, \$10; 3 years, \$22. Single issues, \$3. Address subscription orders, renewals, and changes of address to *Oregon Geology*, Suite 965, 800 NE Oregon Street # 28, Portland 97232. POSTMASTER: Send address changes to *Oregon Geology*, Suite 965, 800 NE Oregon Street # 28, Portland 97232-2162.

Oregon Geology is designed to reach a wide spectrum of readers interested in the geology and mineral industry of Oregon. Manuscript contributions are invited on both technical and general-interest subjects relating to Oregon geology. Two copies of the manuscript should be submitted. If manuscript was prepared on common word-processing equipment, a file copy on diskette should be submitted in place of one paper copy (from Macintosh systems, high-density diskette only). Graphics should be camera ready; photographs should be black-and-white glossies. All figures should be clearly marked; figure captions should be together at the end of the text.

Style is generally that of U.S. Geological Survey publications. (See USGS *Suggestions to Authors*, 7th ed., 1991, or recent issues of *Oregon Geology*.) Bibliography should be limited to references cited. Authors are responsible for the accuracy of the bibliographic references. Include names of reviewers in the acknowledgments.

Authors will receive 20 complimentary copies of the issue containing their contribution. Manuscripts, letters, notices, and meeting announcements should be sent to Beverly F. Vogt, Publications Manager, at the Portland office (address above).

Permission is granted to reprint information contained herein. Credit given to the Oregon Department of Geology and Mineral Industries for compiling this information will be appreciated. Conclusions and opinions presented in articles are those of the authors and are not necessarily endorsed by the Oregon Department of Geology and Mineral Industries.

Cover illustration

Collage of samples from reconstructions of plants and animals from fossils of the John Day Fossil Beds National Monument. Illustrations and discussion of such reconstructions are contained in the article beginning on next page.

In memoriam: Volunteer Shirley O'Dell

Shirley O'Dell, volunteer at the Nature of the Northwest Information Center, died March 30, 1996.

Born in Minneapolis, Minnesota, she spent most of her life in Portland, Oregon. She was a secretary at Charles F. Berg, at an investment company, and at Columbia Grain Company. After her retirement, she worked as a volunteer at St. Vincent's Medical Center Gift Shop and at the Nature of the Northwest Information Center. She was a member of the Geological Society of the Oregon Country (GSOC) for more than 30 years and held various elective and appointive offices, including secretary and auditor.

Shirley was an avid worldwide traveler. She is missed at the Center because she contributed so much as a volunteer—her knowledge about Oregon, interest in geology, and willingness to do careful, meticulous work.

She is survived by one brother and many friends. □

DOGAMI will map tsunami hazard zones at Newport and Seaside-Gearhart

The Oregon Department of Geology and Mineral Industries (DOGAMI) has received support to produce tsunami hazard maps for the Yaquina Bay area of Newport and for the cities of Seaside and Gearhart. The Seaside-Gearhart project is funded by a grant of \$132,000 from the Petroleum Antitrust Grant Program administered by the Oregon Department of Justice. The Newport project is funded by a grant of \$85,000 from the National Earthquake Hazard Reduction Program of the U.S. Geological Survey.

Both projects will produce maps of expected tsunami flooding from undersea earthquakes on the nearby Cascadia subduction zone fault system. Great (magnitude 8–9) earthquakes on this fault system have caused large tsunamis to strike the Oregon coast every 300 to 500 years, the last one being 300 years ago. Such tsunamis strike the coast within 30 minutes or less after the earthquake, so evacuation guided by accurate knowledge of possible flooding zones is essential.

DOGAMI is working in collaboration with the Oregon Graduate Institute of Science and Technology and the Geological Department of Portland State University to generate the necessary data for these maps. For information about the projects, contact George Priest at the Portland office of DOGAMI, phone 503-731-4100. □

DOGAMI now has Internet home pages

The Oregon Department of Geology and Mineral Industries (DOGAMI) can now be found on the Internet with its own home pages:

The general Department address

<http://sarvis.dogami.state.or.us>

The Nature of the Northwest Information Center:

<http://www.naturenw.org>

The Mined Land Reclamation Program:

<http://www.proaxis.com/~dogami/mlrweb.shtml> □

Reconstructions of Eocene and Oligocene plants and animals of central Oregon

by Gregory J. Retallack, Erick A. Bestland, Department of Geological Sciences, University of Oregon, Eugene, OR 97403-1272, and Theodore J. Fremd, John Day Fossil Beds National Monument, HC 82 Box 126, Kimberly, OR 97848-9701

ABSTRACT

Tertiary fossil plants and mammals of the Clarno and Painted Hills units of the John Day Fossil beds have now been placed within a stratigraphic framework created by excavating trenches through poorly exposed parts of the sequence. The middle-late Eocene Nut Beds locality in the Clarno Formation of the Clarno area is at the top of a sequence of volcanic mudflows, that include fossil plant localities near the Hancock Tree and the Fern Quarry of about the same geological age. Reconstructions are presented here of an extinct tree (*Meliosma beusekomii*), a tortoise (*Hadrianus corsoni*) and a variety of fossil mammals (*Telmatherium manteoceras*, *Hyrachyus eximius*, *Orohippus major*, *Patriofelis ferox*) from the Nut Beds, and of an extinct plane tree (*Macginitiea angustiloba*), from near the Hancock Tree. Also reconstructed are late Eocene mammals (*Protitanops curryi*, *Plesiocolopirus hancocki*, *Telestaceras radinskyi*, *Haplohippus texanus*, *Diplobunops crassus*, *Hemipsalodon grandis*) similar to those from the Mammal Quarry of the uppermost Clarno Formation. Newly discovered in the Clarno area but in the lower John Day Formation are fossil plants and mammals of late Eocene age. Early Oligocene lake beds of the middle Big Basin Member of the lower John Day Formation in the Painted Hills include a variety of plants, dominated by an extinct alder (*Alnus heterodonta*), reconstructed here. Lignites stratigraphically higher within the middle Big Basin Member have yielded mostly remains of dawn redwood (*Metasequoia occidentalis*), also reconstructed here. Late Oligocene fossil mammals (*Diceratherium annectens*, *Miohippus quartus*, "Entelodon" *calkinsi*, *Hypertragulus hesperius*, *Eporeodon occidentalis*, *Nimravus brachyops*) from the Turtle Cove Member of the John Day Formation also are reconstructed.

INTRODUCTION

Our impressions of the prehistoric past are based less on illustrations of fossil bones and leaves than on the work of such artists as Charles Knight, Bruce Horsfall, Zdenek Burian, Jay Matternes, Frank Knight, and Peter Murray. Their restorations of past landscapes are more powerful coherent images than fragmentary fossils scattered in museum drawers and published in technical journals. Their artistry also makes obvious testable hypotheses of the nature of past life, hypotheses that can prod the search for deeper understanding. In this spirit, we offer new reconstructions of some fossil plants and animals from the Eocene-Oligocene parts of the Clarno and John Day Formations of central Oregon.

Central Oregon has long been known as an important region for Cenozoic nonmarine fossils now conserved in the John Day Fossil Beds National Monument (Figure 1). Fossil leaves were discovered in the Painted Hills by soldiers using the military road from John Day to The Dalles in the 1860s. Thomas Condon, then a missionary in The Dalles, made the first scientifically useful collections of plant and mammal fossils from the region in 1865 (Clark, 1989) and shipped them east for expert description (Marsh, 1873, 1874, 1875; Newberry, 1898; Knowlton, 1902). Expeditions from the University of California at Berkeley revisited these sites at the turn of the century and discovered plant fossils in the Clarno area (Merriam, 1901a,b, 1906; Sinclair, 1905; Merriam and Sinclair, 1907). The most significant sites in the Clarno area were first developed by amateurs, the Nut Beds by Thomas Bones, beginning in 1943 (Stirton, 1944; R.A. Scott, 1954; Bones, 1979; Manchester, 1994a), and the Mammal Quarry by Lon Hancock in 1955 (Mellest, 1969; Hanson, 1973, 1989; Pratt, 1988). The fossil riches of the area are far from exhausted, and paleontological research is still active (Retallack, 1991a,b; Manchester and Kress, 1993; Manchester, 1994a,b, 1995; Manchester and others, 1994; Meyer and Manchester, unpublished data, 1995; Fremd, 1988, 1991, 1993; Fremd and Wang, 1995). John Day fossils present striking evidence of paleoclimatic and evolutionary change on a 50-million-year time scale (Chaney, 1948a; Ashwill, 1983; Wolfe, 1992; Bestland and others, 1994). Emphasis of this paper is on their diversity and quality of preservation that allows an unusually detailed view of plants and animals on land in the distant geological past.

EOCENE FOSSILS OF THE CLARNO AREA

The geological sequence of fossil localities in the type area of the Clarno Formation near Clarno has remained unclear until recently because of the complicating effects of a local dacite intrusion and poor outcrop of clayey strata in badlands and grassy slopes. We addressed both problems by digging trenches to improve exposure and found, for example, that the dacite intrusion has a capping paleosol, so that it was a volcanic hill overlapped by fossiliferous rocks of the upper Clarno Formation (Bestland and Retallack, 1994a; Bestland and others, 1994, 1995). Such stratigraphic data, together with recent single-crystal $^{40}\text{Ar}/^{39}\text{Ar}$ radiometric dates (by C.C. Swisher for Bestland and Retallack, 1994a), have enabled us to establish the stratigraphic order of fossil localities near Clarno (Figure 2).




Figure 1. Location map for Painted Hills and Clarno Units of the John Day Fossil Beds National Monument, showing distribution of Clarno and John Day Formations and older rocks (after Walker, 1977; Walker and Robinson, 1990).

The best known fossil locality in this area is one known as the Clarno Nut Beds, named for an extraordinary diversity of 173 species of fossil fruits and seeds (Manchester, 1994a), fossil wood, leaf impressions, and insects (R.A. Scott, 1954, 1955, 1956; R.A. Scott and Barghoon, 1956; R.A. Scott and others, 1962; Gregory, 1969; Manchester, 1977, 1979, 1980a,b, 1981, 1983, 1987, 1988, 1991; Manchester and Kress, 1993). The diversity of the flora, its vines, bananas, palms, and other taxa, has been regarded as evidence of multitratal rain forest (Manchester, 1994a). However, associated paleosols are Ultisols (Smith, 1988;

Bestland and Retallack, 1994a) rather than the Oxisols typical under equatorial rain forest or "high evergreen selva" (of Gómez-Pompa, 1973). A better modern analog considering the palms, cycads, tree ferns, oaks, planes, and laurels in the fossil flora may be the "high semi-evergreen montane selva of Lauraceae" of tropical Mexico (Gómez-Pompa, 1973). Vertebrate fossils found in the conglomerates of the Nut Beds include tortoise (*Hadrianus* sp. indet.), crocodylian, titanothere (*Telmatherium* sp. indet.), four-toed horse (*Orohippus major*), extinct rhino (*Hyrachyus eximius*), and lionlike extinct carnivore (*Patriofelis ferox*)

(Hanson, 1973; Pratt, 1988). Such a fauna is compatible with a forested habitat and middle Eocene (Bridgerian) age. Late Eocene (Uintan) is indicated by recent radiometric determinations of 43.6 and 43.7 Ma by Vance (1988), using fission tracks, and of 43.7 Ma by Turrin (for Manchester, 1994a), using single-crystal laser-fusion $^{40}\text{Ar}/^{39}\text{Ar}$ on plagioclase crystals. Most of the fossil mammals listed are known also from the Uintan, and there remain problems with the Uintan-Bridgerian boundary (Prothero and Swisher, 1992; Evanoff and others, 1994). A more precise determination than "Bridgerian-Uintan" will have to await further studies of the Wyoming type section and the Clarno fossils.

The relationship of the Nut Beds to other nearby localities has been problematic (Retallack, 1981). On the basis of our correlation of rhyodacitic ashes, unusually thick lahars, and a distinctive amygdaloidal basalt, we place the Nut Beds at the top of the volcanic mudflows in the Clarno Formation. These thick lahars with few and weakly developed paleosols accumulated rapidly, so that the numerous fossil plant localities now known to underlie the Nut Beds are not much different in geologic age. This idea is supported by fission track dates of 44 Ma for the Muddy Ranch tuff and of 44.3 Ma from a dacite cobble in a lahar above the Fern Quarry (Vance, 1988). The dacite cobble was probably derived from the onlapped local dacite dome, which is older than any known Eocene fossil locality in this area. The Muddy Ranch tuff is within a clayey sequence of paleosols between the conglomerates of the Palisades and the overlying conglomerates of Hancock Canyon, which are capped by the Nut Beds.

A newly discovered locality for lignites is within the lahar sequence, buried within the slope beneath the amygdaloidal basalt, 1 km northeast of the Oregon Museum of Science and Industry (OMSI) Hancock Field Station (Bestland and Retallack, 1994a). The thick, peaty surface horizon, associated fossil logs, and tabular systems of thick carbonaceous fossil roots are evidence of swamp forest dominated by dicotyledonous angiosperms. These are preserved in the paleosols as leaf compressions and include laurels (*Litseaephyllum*




Figure 2. Geological section and fossil assemblages of the upper Clarno Formation in the Clarno Unit of the John Day Fossil Beds National Monument (illustrations adapted from Chaney, 1937; R.A. Scott, 1954; MacGinitie, 1969; Mellett, 1969; Brown, 1975; Manchester, 1981, 1986; Crane and Stockey, 1985; Savage and Long, 1986).

presanguinea, *Cinnamomophyllum* sp. cf. "*Cryptocarya*" *eocenica*) and aguacatilla (*Meliosma* sp. cf. *M. simplicifolia*), as well as fragments of a broad-leaved grass or sedge (*Graminophyllum* sp.) and a fern (*Anemia grandifolia*). The dominance of this swamp by dicots is most like swamps of central America south of Mexico (Breedlove, 1973; Porter, 1973; Hartshorn, 1983). In northern Mexico and the United States, in contrast, swamps are dominated

by conifers such as bald cypress (*Taxodium distichum*) (Best and others, 1984). The dominance of swamps of temperate to subtropical climates by taxodiaceous conifers can be traced back in the fossil record at least into Late Cretaceous times (Retallack, 1994b).

The Fern Quarry also is within the lahar sequence of the upper Clarno Formation. Its fossil plants include mainly ferns (*Acrostichum hesperium*) with lesser horsetails (*Equisetum clarnoi*) and dicots (*Joffrea speirsii*, *Quercus* sp., *Cinnamomophyllum* sp. cf. "*Cryptocarya*" *eocenica*). Ferns and horsetails are in growth position, dicots in leaf litters of very weakly developed volcanoclastic paleosols. This herbaceous and shrubby vegetation may have been early in the ecological succession to colonize volcanic ash (Burnham and Spicer, 1986).

The Hancock Tree is a local landmark in Hancock Canyon and is part of an extensive fossil forest within the lahars of the upper Clarno Formation (Manchester, 1986). It is 39 cm in diameter and shows a 279-cm length of exposed trunk above a concealed basal portion of 115 cm to the base of the lahar and 28 cm to the top of the paleosol in which it is rooted: a total preserved length of 422 cm. The Hancock tree has 59 annual rings as well as another 12 mm too deformed to count in its 39-cm diameter. Six other prone fossil logs within the lahar had diameters of 32, 9, 7, 6, 6, and 3 cm. These were thus tall colonizing forests, intermediate between secondary regrowth and old-growth forest. Fossil leaf litters have been recovered from paleosols supporting the permineralized trunks; they include mainly temperate elements such as sycamore (*Macginitiea angustiloba*), katsura (*Joffrea speirsii*), and alder (*Alnus clarnoensis*), with few tropical elements such as laurel (*Cinnamomophyllum* sp. cf. "*Cryptocarya*" *eocenica*). In contrast, leaf litters of a weakly developed paleosol in conglomerates underlying the Nut Beds to the east yielded mainly tropical elements such as aguacatilla (*Meliosma* sp. cf. *M. simplicifolia*) and magnolia (*Magnolia* sp. cf. *M. leei*) with less common walnut (*Juglans* sp. indet.) and sycamore (*Macginitiea angustiloba*). Similarly, paleosols so weakly developed that bedding has persisted within the Nut Beds include aguacatilla (*Meliosma* sp. cf. *M. simplicifolia*), moonseed (*Diploclisia*), icacina vine (*Goweria dilleri*), magnolia (*Magnolia leei*), laurels (*Litseaphyllum praesanguinea*, *L. praelingue*, *L.* sp. cf. "*Laurophyllum*" *merrilli*, *Cinnamomophyllum* sp. cf. "*Cryptocarya*" *eocenica*), tree fern (*Cyathea pinnata*), horsetail (*Equisetum clarnoi*), walnut (*Juglans* sp.), maple (*Acer clarnoense*), alder (*Alnus clarnoensis*), katsura (*Joffrea speirsii*), and sycamore (*Macginitiea angustiloba*). Again, to the east, very weakly developed paleosols have leaf litters of temperate climatic affinities, including leaves of sycamore (*Macginitiea angustiloba*), katsura (*Joffrea speirsii*), alder (*Alnus clarnoensis*), laurel (*Litseaphyllum praesanguinea*), and horsetail (*Equisetum clarnoi*). These leaf litters of both weakly and very weakly developed paleosols in the two areas represent an ecotone between two distinct kinds of

ancient colonizing forest. *Meliosma* today includes species of colonizing forests (van Beusekom and van de Water, 1989), as do living sycamore (*Platanus*) (Peattie, 1991; Manchester, 1986) and alder (Burger, 1983). Vegetation comparable to the *Macginitiea*-dominated fossil assemblages in the eastern part of the OMSI Hancock Field Station is deciduous tropical forest dominated by *Liquidambar macrophylla* found at elevations of 1,000–2,000 m on volcanoes in tropical Mexico (Gómez-Pompa, 1973). *Meliosma*-dominated assemblages in the western part of the OMSI Hancock Field Station were probably comparable to pioneering forests of lowland evergreen rain forest of Mexico ("selva" of Gómez-Pompa, 1973). Such a reconstruction of deciduous forest on volcanic footslopes to the east and semi-evergreen forest on volcanic toeslopes to the west is compatible with a reconstructed Clarno paleogeography of an andesitic stratocone source for lahars to the east of the study area (White and Robinson, 1992). High-elevation deciduous Eocene plants became increasingly important for the evolution of Oligocene and younger cool-temperate deciduous floras of North America (Wolfe, 1987).

The Mammal Quarry is close to the upper contact of the Clarno Formation with the basal ash-flow tuff (member A) of the John Day Formation, which has now been radiometrically dated at 39.22±0.03 Ma, using single-crystal ⁴⁰Ar/³⁹Ar dating (Bestland and Retallack, 1994a). Thus it includes the oldest known mammalian fauna of the Duchesnean North American Land Mammal "Age" (Lucas, 1992). Very weakly developed paleosols and closely associated fluvial deposits in the quarry have yielded crocodylians, creodont carnivores (*Hemiposalodon grandis*), sabre-tooth cats (Nimravidae gen. et sp. indet.), rodents (gen. et sp. indet.), anthracotheres (*Heptacodon* sp. indet.), oreodons (*Diplobunops* sp. indet.), rhinoceroses (*Teletaceras radinskyi* and cf. "*Procadurcodon*" sp. indet.), tapirs (*Plesiocolopirus hancocki* and *Protapirus* sp. indet.), and horses (*Epihippus gracilis* and *Haplohippus texanus*) (Mellett, 1969; Hanson, 1973, 1989; Schoch, 1989; Lucas, 1992). Some of these creatures, particularly the common small rhinoceros (*Teletaceras radinskyi*), evidently died locally, because they are represented by moderately complete though disarticulated specimens. Their carcasses accumulated on a fluvial point bar as isolated elements or groups of bones still united by flesh (Pratt, 1988).

Also found in the Mammal Quarry was a fossil flora of fruits and seeds (McKee, 1970; Manchester, 1994a), including tropical vines (*Odontocaryoidea nodulosa*, *Diploclisia* sp. indet., *Eohypserpa* sp. indet., *Iodes* sp. indet., *Iodicarpa* sp. indet., *Palaeophytocrene* sp. cf. *P. foveolata*, *Ampelocissus* sp. indet., *Vitis* sp. indet., *Tetrastigma* sp.), sycamores (*Platananthus synandrus*), dogwoods (*Mastixioidiocarpum oregonense*), alangiums (*Alangium* sp. indet.), cashews (*Pentoperculum minimus*), and walnuts (*Juglans clarnensis*). Late Eocene mammals here evidently lived in forests, but Luvisol paleosols in association with these fossils are evidence for a cooler and drier climatic

regime than during deposition of the Nut Beds (Bestland and Retallack, 1994a).

New fossil localities were also discovered within the lower John Day Formation in footslopes of Iron Mountain, stratigraphically below a distinctive thick tuff radiometrically dated (using the single-crystal laser-fusion $^{40}\text{Ar}/^{39}\text{Ar}$ technique) at 38.4 ± 0.07 Ma (C. Swisher for Bestland and Retallack, 1994a). This is still within the late Eocene and Duchesnean (Swisher and Prothero, 1990). From a red paleosol described by Getahun and Retallack (1991) immediately below the dated tuff came a fragment of a tusk, probably of the extinct family of hoglike Entelodontidae, and a molar of the small Clarno rhino *Tele-taceras radinskyi*. Brown lake beds with fish scales some 5 m below the tuff have yielded winged fruits of walnutlike plants (*Cruciptera simsoni*, *Palaecarya clarnensis*), fruits of eucommia (*Eucommia* sp. indet.), extinct malvacean flowers (*Florisantia ashwillii*), seeds of cigar-tree (*Catalpa* sp. indet.), and leaves of elm (*Ulmus* sp. indet.), Oregon grape (*Mahonia* sp. indet.), and dawn redwood (*Metasequoia occidentalis*) (Bestland and Retallack, 1994a; S.M. Manchester, oral commun., 1995). This flora and associated Luvisol paleosols are evidence for forests of lesser stature, more open and deciduous than earlier in the Eocene.

The "Slanting Leaf Beds" is a well-known locality higher within the John Day Formation in the Clarno area and now radiometrically dated at 33.62 ± 0.19 Ma (Swisher for Bestland and Retallack, 1994a), which is very early in the Oligocene (Swisher and Prothero, 1990). These tuffaceous lacustrine beds have yielded a very different fossil flora dominated by alder (*Alnus heterodonta*) and dawn redwood (*Metasequoia occidentalis*). This is one of the many Bridge Creek floras especially well known near Fossil and in the Painted Hills (Manchester and Meyer, 1987; Kvacek and others, 1991; Manchester, 1992; Meyer and Manchester, unpublished data, 1995). The dramatic floral change from floras of tropical to those of temperate climatic affinities is re-




Figure 3. Geological section and fossil assemblages of the lower John Day Formation in the Painted Hills Unit of the John Day Fossil Beds National Monument (illustrations adapted from Knowlton, 1902; Sinclair, 1905; Osborn, 1918; Peterson, 1920; Toohey, 1959; Lander, 1977; Tanai and Wolfe, 1977; Wolfe and Tanai, 1987; Meyer and Manchester, unpublished data, 1995).

flected in a change in paleosols from red Luvisols to a mix of Inceptisols, Vertisols and Luvisols (Bestland and Retallack, 1994a), also seen in the Painted Hills.

OLIGOCENE FOSSILS OF THE PAINTED HILLS

The principal fossil resources protected by the Painted Hills Unit of the John Day Fossil Beds are the lake beds with leaves of the Bridge Creek flora (Figure 3), first col-

lected by Thomas Condon in 1865 for Newberry (1898) and Knowlton (1902) and then made famous by pioneering papers in quantitative paleoecology by Chaney (1924, 1925a). Dominance of these shales by plants of temperate climatic affinities such as oak (*Quercus consimilis*), alder (*Alnus heterodonta*), and dawn redwood (*Metasequoia occidentalis*) is in striking contrast to the tropical affinities of fossil plants in the Clarno Formation (Chaney, 1948a). At first, Chaney (1924, 1938) compared the assemblage with that of redwood forests of California. When it was discovered in 1941 that dawn redwood was still living in China, Asian mixed mesophytic forests became a better model (Chaney, 1948a,b, 1952, Wolfe, 1981, 1992; Manchester and Meyer, 1987). In contrast, our new collections indicate that *Metasequoia* was a swampland tree.

One problem with leaves in lake beds is the way in which elements of different plant communities are mixed with and overwhelmed by deciduous lake-margin species. We have been able to demonstrate such biases in the Bridge Creek flora with new collections of plants from leaf litters of paleosols associated with the lake beds (Bestland and Retallack, 1994b). A small fossil flora of leaves found in the surface horizon of a weakly developed green paleosol underlying the lake beds included dawn redwood (*Metasequoia occidentalis*), nutmeg tree (*Torreya* sp. indet.), grass (*Graminophyllum* sp. indet.), laurel (*Cinnamomophyllum bendirei*), and walnut (*Juglans cryptata*). Remains of walnuts were most common, followed by the laurels. The dawn redwood is represented by only one foliar spur. The paleosol and plants combined indicate semi-evergreen mesophytic forest of bottomlands. A very different assemblage was found in a manganese-stained shale overlying a very weakly developed paleosol within the lake beds: mainly leaves of alder (*Alnus heterodonta*) with oak (*Quercus consimilis*) and yellowwood (*Cladrastis oregonensis*). Alders in particular are well-known pioneer plants of disturbed ground, both in tropical and temperate regions (Burger, 1983), and this interpretation is supported by the minimal development of the fossil soil. Lignitic paleosols overlie a pumice-charcoal tuff above the lake beds. These ancient soils of swamps are overwhelmingly dominated by dawn redwood (*Metasequoia occidentalis*), associated with rare bracken fern (*Pteridium calabazensis*), grass (*Graminophyllum* sp. indet.), alder (*Alnus heterodonta*), and extinct basswood (*Plafkeria obliquifolia*). These paleosol assemblages indicate that the Bridge Creek flora of lacustrine shales is a mix of at least three distinct plant communities: semi-evergreen lowland laurel forest, dawn redwood swamp, and deciduous alder pole woodlands.

The age of the Bridge Creek flora in the Painted Hills is constrained by new single-crystal laser-fusion $^{40}\text{Ar}/^{39}\text{Ar}$ radiometric dates of 32.99 ± 0.11 Ma for a biotite tuff below the lake beds and 32.66 ± 0.03 Ma for tuffs above the lake beds and lignites, both within the middle Big Basin Member of the John Day Formation (C.C. Swisher for Bestland and Retallack, 1994b). Like the "Slanting Leaf Beds" of the

Clarno area, these also are earliest Oligocene in age. At some time between 33 Ma and the 38-Ma fossil flora from the Clarno area discussed above, there was a climatic deterioration from mean annual precipitation of 1,500 mm to 1,000 mm (Leopold and others, 1992), from mean annual temperature of 20°C to 12°C , and from mean annual range of temperature of 10°C to 25°C (Wolfe, 1992). A comparable deterioration to temperate climate is indicated by the transition from kaolinitic and iron-rich, red, lateritic Ultisol and Luvisol paleosols of the lower Big Basin Member of the John Day Formation, which are erosionally truncated and abruptly overlain by smectitic and tuffaceous Inceptisol, Vertisol, and Luvisol paleosols of the middle Big Basin Member (Bestland and others, 1994, 1995, 1996). The erosional contact probably corresponds to the Eocene-Oligocene boundary now recognized at 34 Ma (Swisher and Prothero, 1990), judging from radiometric dates from underlying units: a new $^{40}\text{Ar}/^{39}\text{Ar}$ single-crystal radiometric date of 39.72 ± 0.03 Ma for the basal ash-flow tuff of the John Day Formation (C.C. Swisher for Bestland and Retallack, 1994b) and a date of 37 Ma for the andesite of Sand Mountain (Hay, 1963; Evernden and James, 1964). Our correlation of these units and assessment of the degree of development of lateritic paleosols of the lower John Day Formation suggest that the Eocene-Oligocene climatic deterioration was abrupt (Retallack, 1992; Rice, 1994; Bestland and others, 1996).

The badlands of Carroll Rim in the Painted Hills and the nearby footslopes of Sutton Mountain have yielded only sparse and fragmentary mammal fossils from the Turtle Cove Member below the massive ash-flow tuff of Carroll Rim ("Picture Gorge tuff" or Member H of Robinson and others, 1984). The Turtle Cove Member here is dated by new determinations of 29.75 ± 0.02 , 29.70 ± 0.06 , 28.65 ± 0.05 , and 28.65 ± 0.07 Ma on tuffs and the cliff-forming ash flow (last two dates), respectively (C.C. Swisher for Bestland and Retallack, 1994b). Comparable late Oligocene (early Arikareean) fossil mammals are more abundant and better known from the Turtle Cove Member in the John Day Valley south of Kimberly (Fremd, 1988).

Fossil mammal fragments were mainly found in paleosols and can be grouped by paleosol type. Some of these have brown, finely structured, near-mollic surface horizons over a subsurface horizon of calcareous nodules, as in soils now supporting grassy woodland and wooded grassland. Fossil mammals in these paleosols include mouse deer (*Hypertragulus hesperius*), oreodonts (Merycoidontidae gen. et sp. indet.), and rhinoceros (*Diceratherium* sp. cf. *D. annectens*). Other paleosols are comparable in their brown, finely structured surface horizons, but lack nodules, which indicates a shorter time for formation but comparable vegetation. Mammals of these paleosols include an unidentifiable mammalian navicular, a distal phalanx of an oreodont (Merycoidontidae), and a cuboid similar to that of a giant, hoglike entelodon (cf. *Entelodon* sp. indet.). Very weakly developed paleosols probably supported streamside grassy

vegetation early in ecological succession to communities represented by the other paleosols. Mammals of these transient communities include extinct squirrel-like rodent (Eutypomyidae gen. et sp. indet.), oreodon (*Agriochœrus* sp. indet., *Eporeodon* sp. indet.), three-toed horse (*Miohippus* sp. cf. *M. quartus*), and two-horned rhinoceros (*Diceratherium* sp. cf. *D. annectens*). Other paleosols are green and clayey with subsurface calcareous nodules and probably supported seasonally wet alluvial grassy woodland. Comparable paleosols near Foree to the east have yielded oreodont (*Eporeodon occidentalis*), three-toed horse (*Miohippus* sp. cf. *M. anceps*), and foxlike dog (*Mesocyon coryphaeus*). No red, clayey paleosols were seen in the Turtle Cove Member of the Painted Hills, but red paleosols in Rudio Creek canyon near Kimberly yielded three-toed horse (*Miohippus equiceps*). Although endocarps of hackberry (*Celtis willistoni*) (Cockerell) Berry 1928 (Chaney 1925b) are the only plant fossils found in the Turtle Cove Member, paleosols indicate a mosaic of woodland in hills and footslopes, wooded grassland and grassy woodland in valley bottoms, and local grassy woodlands in seasonally wet bottomlands. Mammalian collections from particular paleosols are as yet inadequate to demonstrate their ecological preferences. Nevertheless, the paleoenvironmental reconstruction from paleosols is compatible with the mix of arboreal squirrels, browsing horses and oreodonts, and grazing rhinos, similar to assemblages in the Great Plains regarded as adapted to grassy woodland and wooded grassland (Webb, 1977; MacFadden, 1992; Prothero and others, 1996).

Paleosols of the Turtle Cove Member indicate much drier and probably also cooler paleoclimatic conditions than earlier in the Oligocene. Depth to the calcareous nodules is strongly correlated with mean annual rainfall in modern soils (Retallack, 1994a), and application of these relationships with allowances for compaction of paleosols in Carroll Rim gives mean annual rainfall of 614±141 mm to 911±141 mm (Bestland and Retallack, 1994b). This range of about 500–1,000 mm for the Turtle Cove Member can be compared to 800–1,000 mm for the upper Big Basin Member and 1,000–1,200 mm for the middle Big Basin Member, derived by comparison with comparable soils in Mexico (Bestland and Retallack, 1994b). Paleoclimatic drying continued into the early Miocene, judging from our reconnaissance observations of the upper John Day Formation on Sutton Mountain. The John Day Formation preserves a long-term paleoclimatic record that complements and correlates well with the geological record of global change in deep-sea cores (Bestland and Retallack, 1994b).

RECONSTRUCTED FOSSIL PLANTS

An outstanding advantage of preservation of plants in paleosols is the lack of mixing and separation of different parts of different plants. The leaves, fruits, flowers, and wood are preserved near their place of growth, little mixed by transport in streams or drifting off into lake beds. Such

fossiliferous paleosols are known in the lower part of the Clarno Nut Beds and within the lignites and lake beds of the Painted Hills, and they make easier the difficult task of reconstructing ancient plants from a jumble of different plant remains (Manchester, 1981). Physical attachment of one part of a plant to another, such as fruits on leafy twigs, is the most compelling, but unfortunately rare, evidence for reconstructing fossil plants. Comparison with modern plants is also useful, particularly if there is only one fossil species of fruit, wood, flower, and leaf comparable to a particular modern plant. In this way, it is hoped to gain a complete concept of particular ancient plants and their evolution. As this work proceeds, it is revealing that some parts of plants such as fruits and seeds evolve at rates different from others such as wood and leaves (Manchester, 1986, 1987; Manchester and Wheeler, 1993). This mosaic evolution not only explains differing interpretations of ancient vegetation based on wood, fruits, and leaves but is of interest for understanding the evolutionary process itself.

Meliosma beusekomii

Meliosma beusekomii Manchester (1994a) is a middle Eocene fossil species named for fruits common in the Clarno Nut Beds of the Clarno Formation (Figure 4). The fruits are drupes, but the preserved parts are characteristically dimpled stones or endocarps (Manchester, 1994a). *Meliosma beusekomii* endocarps are very similar to those of living species such as *M. pinnata* and *M. simplicifolia*. Leaves comparable to *M. simplicifolia* also are common in the Nut Beds and in other fossil leaf litter assemblages of the upper Clarno Formation (Manchester, 1980b). Our reconstruction of the shoot with fruits and leaves is based on that of *M. simplicifolia* (van Beusekom and van de Water, 1989). Wood of *Meliosma* has also been found in the Nut Beds (Manchester, 1980b) and was sketched from photomicrographs generously supplied by Steven Manchester.

The large leaves (up to 21 cm long) of this extinct Eocene species of *Meliosma* have a pronounced drip tip, as is typical for plants of tropical rainy climates. *Meliosma* today is widespread in the humid tropics and subtropics of eastern Asia, Indonesia, Central America, and the West Indies (Gentry, 1980; van Beusekom and van de Water, 1989). *Meliosma* includes both shrubs and trees, and its family Sabiaceae includes vines (Gentry, 1980). Permineralized fossil wood of *Meliosma* is known only as waterworn fragments but is evidence that *M. beusekomii* was a tree. It probably was not a large tree, because its leaves dominate the litters of very weakly developed paleosols that lack large root traces or differentiated soil horizons. These paleosols in the Nut Beds and elsewhere are associated with fluvial facies (Retallack, 1991a), so that *M. beusekomii* can be envisaged a pole tree of woodlands early in ecological succession of recovery from stream flooding.

Macginitiea angustiloba

Macginitiea angustiloba (Lesquereux) Manchester




Figure 4. Reconstruction of a fossil aguacatilla (*Meliosma beusekomii*) from the middle-late Eocene "Nut Beds" of the Clarno Formation at Hancock Field Station (data from Bones, 1979; Manchester, 1981, 1994a; van Beusekom and van de Water, 1989).

(1986) is the name of a palmately lobed leaf common in lahars of Hancock Canyon and elsewhere in the Eocene Clarno Formation (Figure 5). The leaf is similar to those of living sycamores and planes (*Platanus*). The Hancock tree is a well-known local landmark in Hancock Canyon and is a katsura (*Joffrea speirsii*) (Retallack, 1991a), but the other

common species within the same paleosol is *Macginitiea angustiloba*. Stumps of *Macginitiea angustiloba* are recognizable in hand specimen by their characteristic wide rays, and this wood has been referred to *Plataninium haydenii* (Wheeler and others, 1977; R.A. Scott and Wheeler, 1982). The globose pistillate and staminate heads and isolated anthers of this plant have also been given separate names, *Macginicarpa glabra*, *Platananthus synandrus*, and *Macginistemon mikanooides*, respectively (Manchester, 1986), in the widely used organ-generic procedure of paleobotany. There is little doubt from their joint association in numerous localities, including lake beds and fluvial sandstones as well as the paleosols of Hancock Canyon and the Nut Beds, that these represent parts of the same extinct species of sycamore-like plant (Manchester, 1986).

Like modern sycamores, *Macginitiea angustiloba* had leaves with abscission scars, indicating it was deciduous. Its separate pistillate and staminate flowering heads produced abundant small (16–18 μ) pollen and small (6–8 mm), hard fruits, which are indications that it was pollinated and dispersed by wind and water (Manchester, 1986). *Macginitiea angustiloba* had slightly smaller pollen, better developed perianth, and more consistently pentamerous flowers than modern sycamores, which may indicate facultative insect pollination or an evolutionary legacy of insect pollination. *Macginitiea angustiloba* also lacked hairs that facilitate wind dispersal of fruits of living sycamores. This combination of largely wind and water pollination and dispersal with some hints of facultative animal involvement are common in plants that colonize disturbed ground. Like living sycamores (*Platanus*), many comparable fossils as old as Cretaceous also were trees of disturbed streambanks (Retallack and

Dilcher, 1981). Such an ecology for the Clarno plant is indicated in Hancock Canyon, where trunks attributed to *Macginitiea angustiloba* are rooted in a weakly developed paleosol capping fluvial deposits interbedded with lahars (Bestland and Retallack, 1994a). *Macginitiea angustiloba* is the most prominent of several plants in the Clarno For-

mation that were deciduous, wind and water pollinated and dispersed, and with cool-temperate climatic affinities. Unlike diverse assemblages of the Nut Beds, low diversity fossil forests of Hancock Canyon may represent a flora of higher elevations on the volcanic cone, like the *Liquidambar* forests of San Martin Volcano in Veracruz, Mexico (Gómez-Pompa, 1973).

Metasequoia occidentalis

Foliage, cones, and seeds of the dawn redwood *Metasequoia occidentalis* (Newberry) Chaney (1951) are common in the Bridge Creek flora of the middle Big Basin Member of the John Day Formation at many localities, including the Painted Hills, the Slanting Leaf Beds of the Clarno area, and the well-known bank behind the High School in Fossil (Manchester and Meyer, 1987). In the Painted Hills, foliage of this plant is found in nearly monospecific assemblages with permineralized coniferous wood in a sequence of lignites. Exceptionally well preserved permineralized wood, cones, and foliage of *Metasequoia* also are known from the middle Eocene Princeton chert of British Columbia (Rothwell and Basinger, 1979; Basinger, 1981, 1984), which is a permineralized peat swamp. Although these Eocene fossils have been referred to the species *Metasequoia milleri*, they are externally identical to living dawn redwood *M. glyptostroboides*, which cannot be readily distinguished from the fossil species for leaf impressions, *M. occidentalis*. *Metasequoia milleri* differs from *M. glyptostroboides* in its higher wood rays and two additional resin canals in the distal portion of the microsporophyll of the pollen cone (Rothwell and Basinger, 1979; Basinger, 1981). Plants of peat swamps are known to evolve very slowly; for example, many species known from permineralized remains range through the entire 30 Ma of the Pennsylvanian with little evolutionary change (DiMichele and others, 1987). Similarly, *M. occidentalis*, *M. milleri*, and *M. glyptostroboides* may all repre-




Figure 5. Reconstruction of a fossil plant (*Macginitiea angustiloba*) from the middle-late Eocene conglomerates of Hancock Canyon of the Clarno Formation at Hancock Field Station, Oregon (data from Wheeler and others, 1977; R.A. Scott and Wheeler, 1982; Manchester, 1986).

sent the same long-lived species. *Metasequoia* has long been known as a living fossil, because it was discovered alive in China after description as a fossil (Chaney, 1948a). This raises the interesting procedural problem of a living genus with a fossil type specimen (Schopf, 1948).

Metasequoia has been extinct in North America since




Figure 6. Reconstruction of a fossil dawn redwood (*Metasequoia occidentalis*) from the early Oligocene middle Big Basin Member of the John Day Formation in the Painted Hills, Oregon (data from Merrill, 1948; Manchester and Meyer, 1987; Meyer and Manchester, unpublished data, 1995; additional details from the probably conspecific *M. milleri* of Rothwell and Basinger, 1979; Basinger, 1981, 1984).

late Miocene time (Chaney, 1948a), and the geologically youngest fossils found anywhere are Pliocene (Miki, 1941). Several hundred *Metasequoia* trees now growing in Oregon are descended from seeds distributed by Ralph Chaney in 1950. During his visit to China in the spring of 1948, he

was unable to obtain seeds, but he distributed seeds purchased from China by E.D. Merrill (1948) of Harvard's Arnold Arboretum. A particularly fine tree grown from Chaney's original distribution grows near Cascade Hall on the University of Oregon campus. It stands under the gaze of a bronze bust of Thomas Condon, who first discovered these Oregon fossils. Sketches of this tree were used for our reconstruction (Figure 6).

Metasequoia is seasonally deciduous like larch (*Larix*) and differs in this respect from most conifers, which are evergreen. The pollen cones are formed with the soft, light-green spring leaves, and the seed cones do not mature until late in the fall. Like most conifers it is wind pollinated, and seed is dispersed by wind and water. *Metasequoia occidentalis* has long been regarded as a dominant of a mixed conifer-angiosperm flora (Chaney, 1924, 1925a; Wolfe, 1981, 1992), and this has been regarded as confirmed by its occurrence in China (Chaney, 1948a, Chu and Cooper, 1950; Hu, 1980). However, this small area of northwest Hupei Province has been greatly disturbed by agriculture. *Metasequoia glyptostroboides* was found in ravines and embankments around paddy fields that have submerged at least two hundred trunks of a relatively pure stand of these plants (Bartholomew and others, 1983). Its common occurrence with angiosperm leaves in the various Bridge Creek floras could be a result of mixing in lake deposits. A new view of the natural habitat of *Metasequoia* emerges from its dominance in the lignites of the Painted Hills (Bestland and Retallick, 1994b) and permineralized peats of British Columbia (Rothwell and Basinger, 1979, Basinger, 1981, 1984). *Metasequoia occidentalis* and the likely conspecific *M. milleri* can now be seen as swamp cypresses

comparable with living bald cypress (*Taxodium distichum*) of Okefenokee Swamp and the Florida Everglades (Best and others, 1984). Both *Taxodium* and *Metasequoia* are in the family Taxodiaceae, and Newberry (1883) may not have been far off the mark when he included fossils now called

Metasequoia in *Taxodium*. These modern swamps have a low diversity complement of angiosperms, and some angiosperm fossils also have been found with *Metasequoia* in the fossil peats of the Painted Hills (Bestland and Retallack, 1994b) and British Columbia (Stockey, 1987; Cevallos-Ferriz and Stockey, 1988a,b, 1989, Erwin and Stockey, 1989a,b, 1994; 1990a,b, 1991; Stockey and Pigg, 1991; Cevallos-Ferriz and others, 1993). However, these cannot be taken as representative of forests of well-drained soils away from the swamps. Much has been made of the climate of relict populations of *Metasequoia* in Hupei to infer Oligocene paleoclimate of Oregon (Chaney, 1948a). As a swamp plant, however, its paleoclimatic significance is limited, because it lived in soils isolated from regional climate by local waterlogging. Its temperature tolerances in the past were probably also cool-temperate to subtropical, like swamp cypresses today, which dominate swamps as far south as northern Mexico but are replaced by angiosperm swamps in the Central American tropics (Breedlove, 1973; Porter, 1973, Hartshorn, 1983).

Alnus heterodonta

Alder leaves and catkins are among the most common fossils in Bridge Creek floras of the middle Big Basin Member of the John Day Formation (Manchester and Meyer, 1987; Meyer and Manchester, unpublished data, 1995). Several species of alder have been recognized, even within single localities (Klucking, 1956), but *Alnus heterodonta* is the most common leaf in the Painted Hills (Figure 7). Young leaves, seeds, and staminate and fruiting catkins also are found occasionally (Meyer and Manchester, unpublished data, 1995). The attribution of all these remains to one plant is based on association and comparison with living alder.

The inconspicuous flowers, small, flattened seeds, and leaf abscission scars of the fossils are evidence that this extinct species was wind pollinated and dispersed and seasonally deciduous like living alder and others of the family Be-




Figure 7. Reconstruction of a fossil alder (*Alnus heterodonta*) from the early Oligocene middle Big Basin Member of the John Day Formation in the Painted Hills, Oregon (data from Klucking, 1956; Wood, 1974; Manchester and Meyer, 1987; Meyer and Manchester, unpublished data, 1995).

tulaceae in cool-temperate climatic regions of the northern hemisphere. Because these leaves dominate the leaf litter of a very weakly developed paleosol in the lake beds of the Painted Hills (Bestland and Retallack, 1994b), it is likely that they were plants of pole woodlands early in the ecological succession to recover disturbed ground. Such an oppor-




Figure 8. Reconstructions of middle-late Eocene (Bridgerian-Uintan) animals from the Nut Beds of the Clarno Formation at the OMSI Hancock Field Station, Oregon (A and scale from R.B. Horsfall for W.B. Scott, 1913; B from E.M. Fulda for Osborn, 1929; E from C.E. Knight for Osborn, 1910; others original with data from Hay, 1908).

tunistic ecology is characteristic of many alder species (Burger, 1983), including the red alder (*Alnus rubra*) of Oregon, which forms roadside and streamside pole woodlands, and extensive stands for some 50–80 years after clear-cutting or fire (Peattie, 1991). Neither the swamp-bound *Metasequoia occidentalis* nor the ephemeral *Alnus heterodonta* are likely to have been abundant in well-drained soils during Oligocene time, despite their dominance of lacustrine leaf accumulations.

RECONSTRUCTED FOSSIL ANIMALS

Complete skeletons or natural mummies of extinct animals are ideal for reconstructing their appearance and lifestyle (Savage and Long, 1986). Few of these have been found in the John Day Fossil Beds, where fleshing out the fragments must rely on more complete skeletons from other

parts of North America and inferences from comparable modern mammals. Fortunately, there is a long tradition of fossil vertebrate reconstruction (Osborn, 1910; W.B. Scott, 1913). More than ever before, we have the resources to conjure up visions of extinct beasts from isolated teeth and bones.

Hadrianus corsoni

Only a fragment of this thick-shelled tortoise has been found in the middle Eocene Clarno Nut Beds, and it is not referable to a particular species. The reconstruction offered here (Figure 8C) is *Hadrianus corsoni* (Leidy) Cope (1873), represented by complete carapaces and limb bones from middle Eocene (Bridgerian) rocks south of Fort Bridger, Wyoming (Hay, 1908). The genus *Hadrianus* ranges through much of the Eocene (Wasatchian to Uintan)

(Hutchinson, 1992). These were land tortoises of the family Testudinidae, which are all herbivores and include the well-known Galapagos tortoises (*Geochelone*) (Carroll, 1988).

Telmatherium manteoceras

Molars of this brontothere are the most common vertebrate fossils in the middle Eocene Clarno Nut Beds and may represent a new species of *Telmatherium*. Our reconstruction (Figure 8B) is based on one by E.M. Fulda (for Osborn, 1929) of *Telmatherium manteoceras* (Hay) Mader (1989), known from good skulls and partial skeletons of middle Eocene (Bridgerian-Uintan) age in the Washakie and Bridger Basins of Wyoming (Osborn, 1929; Eaton, 1985). Painted reconstructions by Charles Knight (for Osborn, 1910, 1929) are very similar to this sketch. Brontotheres (also called titanotheres) are extinct perissodactyls of North America and Asia and represent a distinct evolutionary lineage from rhinoceroses which they vaguely resemble. *Telmatherium* had the low-crowned teeth of a browser, like most bronthotheres. It was small for this group, which evolved from dog-sized animals in the early Eocene to massive rhino-sized animals in the latest Eocene (Carroll, 1988).

Hyrachyus eximius

A variety of skull fragments and teeth in the Clarno Nut Beds belong to *Hyrachyus eximius* Leidy (1871), well known from relatively complete skeletons in middle to late Eocene (Bridgerian to Uintan) deposits of the Bridger and Washakie Basins of Wyoming (W.B. Scott, 1913). Our reconstruction (Figure 8A) is based on the sketch of Bruce Horsfall (for W.B. Scott, 1913), which we prefer to more stocky reconstructions by Zdenek Burian (for Spinar and Burian, 1972), Jay Matternes (for Smithsonian Institution mural, see Dott and Batten, 1981), Michael Long (for Savage and Long, 1986), and Graham Allen (for Dixon and others, 1992). Its low-crowned teeth are those of a browser, but it had padded feet like those of a tapir rather than hooves like modern ungulates. Primitive features of *Hyrachyus* include four toes on the rear limbs and prominent canine teeth. It belongs to an extinct group of mammals (Helatetidae or Hyrachyidae) allied to both tapirs and rhinos and possibly ancestral to both.

Orohippus major

The small, primitive horse *Orohippus major* Marsh (1874) is represented by several teeth and jaw fragments from the Clarno Nut Beds, but good skeletons have been found in the middle Eocene (Bridgerian) of the Bridger Basin of Wyoming (Kitts, 1957). Our reconstruction (Figure 8D) is based on one by Charles Knight (for Osborn, 1910), although others have been done by Zdenek Burian (for Spinar and Burian, 1972) and Jay Matternes (for Smithsonian Institution mural, see Dott and Batten, 1981). With its low-crowned teeth of a browser and four toes on the front limbs and three toes in the rear, this very small

horse is far removed from our modern concept of a horse (MacFadden, 1992). These early forest-dwelling ancestors of modern Equidae lack exact modern analogs.

Patriofelis ferox

Patriofelis ferox (Marsh) Marsh (1878) is represented by a single jaw fragment from the middle Eocene Clarno Nut Beds. A complete skeleton is known from the middle Eocene (Bridgerian) of the Bridger Basin of Wyoming, and this was the basis for our reconstruction (Figure 8E), which is close to that of Charles Knight (for Osborn, 1910). More gracile reconstructions have been made by Bruce Horsfall (for W.B. Scott, 1913), Michael Long (Savage and Long, 1986), and William Scheele (1955). Although it had the appearance of a big cat, *Patriofelis ferox* lacked the precisely bladed carnassial teeth of modern carnivores and belonged to an extinct group of catlike carnivores (Oxyaenidae, Creodonta) (Carroll, 1988).

Protitanops curryi

A variety of limbs and skull fragments of a large brontothere have been found in the late Eocene (Duchesnean) Mammal Quarry of the upper Clarno Formation. Although referred to *Protitanops* (Lucas, 1992), there is doubt about this generic assignment (Mader, 1989). Our reconstruction (Figure 9F) is based on the generally similar *Protitanops curryi* Stock (1936) from the late Eocene (Chadronian) Titus Canyon Formation, east of Death Valley, California. Like *Telmatherium* (Figure 8B), *Protitanops* was a brontothere and a browser. It was much larger than *Telmatherium* and probably had a significant impact in destroying trees and other browse—like modern megaherbivores such as elephants and rhinos (Owen-Smith, 1988). The horns of this and geologically younger brontotheres were probably sexual display organs of male animals.

Plesiocolopirus hancocki

Several skull fragments and teeth of *Plesiocolopirus hancocki* (Radinsky) Schoch (1989) have been found in the Mammal Quarry of the upper Clarno Formation. Radinsky (1963) envisaged this species as intermediate between the earlier *Helalestes nanus* from the middle Eocene (Bridgerian) of Wyoming and the later *Protapirus* from the early Oligocene (Orellan and Whitneyan) Brule Formation of South Dakota (W.B. Scott, 1941)—which he regarded as the first true tapir. Our reconstruction (Figure 9C) was influenced by reconstructions of *Helalestes nanus* by Jay Matternes (see Dott and Batten, 1981) and Bruce Horsfall (for W.B. Scott, 1913) and by those of early Miocene *Miotapirus* by Graham Allen (for Dixon and others, 1992). The Clarno fossils have a large nasal incision and reduced nasal bones. These bones are not so modified as in living tapirs, so the proboscis of *P. hancocki* may not have been so large and flexible. Tapirs are forest browsers, and the teeth of *P. hancocki* are compatible with a similar diet.




Figure 9. Reconstructions of late Eocene (Duchesnean) animals from the Mammal Quarry of the upper Clarno Formation at the OMSI Hancock Field Station, Oregon (scale from R.B. Horsfall for W.B. Scott, 1913; others original with inspiration from Stock, 1936; Russell, 1938; W.B. Scott, 1945; McGrew, 1953; Radinsky, 1963; Mellett, 1969; Savage and Long, 1986; Hanson, 1989; Dixon and others, 1992).

Teletaceras radinskyi

Teletaceras radinskyi Hanson (1989) was a small rhino, represented by skulls and a variety of other bones from the Mammal Quarry of the upper Clarno Formation. Hanson (1989) compared it with *Trigonias osborni* from the Late Eocene (Chadronian) Chadron Formation of South Dakota, a species reconstructed by Irving Biehn (for W.B. Scott, 1913) and Graham Allen (for Dixon and others, 1992). Compared with this geologically younger rhino, the Duchesnean *Teletaceras radinskyi* was a smaller and more gracile animal (Figure 9A). The Clarno rhino may have evolved from immigrant *Teletaceras* from Asia and represents an early true rhino (Rhinocerotidae). As with most rhinos, its teeth are those of a browser.

Haplohippus texanus

The primitive horse *Haplohippus texanus* McGrew, 1953, is represented by a variety of skull fragments and teeth from the Mammal Quarry of the upper Clarno Formation, as well as the late Eocene (Duchesnean) Porvenir local fauna of the Vieja Formation of Texas. The teeth of *Haplohippus texanus* ally it with primitive, forest-browsing horses such as *Orohippus* (Figure 8D), from which it differed by being larger, yet more gracile (Figure 9E). Another horse found in the Clarno Mammal Quarry, *Epihippus gracilis*, on the other hand, is even more gracile and cursorial and has more complex and larger molars, perhaps reflecting a preference for slightly more open habitats. *Haplohippus* was the last of the primitive hyracotheriine horses, but *Epihippus* was in the lineage giving rise to later three-toed horses (McGrew, 1953). Although horse evolution has been portrayed as a unidirectional trend, this and other examples indicate a luxuriant bush of evolutionary branches (MacFadden, 1992).

Diplobunops crassus

A skull of *Diplobunops* from the Mammal Quarry of the upper Clarno Formation is crushed and extensively cracked and may belong to a new species. Our reconstruction (Figure 9D) is based on better preserved material of *Diplobunops crassus* W.B. Scott (1945) from the late Eocene (Uintan) of the Uinta Basin, Utah, and on reconstructions of the allied genus *Agriochoerus* by Bruce Horsfall (for W.B. Scott, 1913). *Diplobunops* and *Agriochoerus* were part of a now-extinct group of creatures grouped into the family Agriochoeridae, which were primitive artiodactyls similar to the extinct but more familiar oreodons common in Oligocene and Miocene parts of the John Day Formation (Carroll, 1988). Their prominent canines and postcranial skeleton are superficially similar to those of dogs, although the tarsus is that of an artiodactyl. The cheek teeth, however, are low crowned and those of a browser.

Hemipsalodon grandis

The large carnivore *Hemipsalodon grandis* (Cope) Russell, 1938, is represented by a crushed but complete skull and lower jaws from the Mammal Quarry of the upper Clarno Formation (Mellett, 1969). This species is known from a variety of localities of late Eocene (Duchesnean and Chadronian) age from Saskatchewan in the north to near the Mexican border of Texas but remains very incompletely known. The bearlike reconstruction offered here (Figure 9B) is based partly on a facial reconstruction by Russell (1938) from Canadian specimens and on an associated astragalus of a plantigrade animal (Mellett, 1969). Although perhaps frugivorous to carnivorous like bears, *Hemipsalodon* was not closely related to them but rather belongs in an extinct group of creodont carnivores (Hyaenodontidae) (Carroll, 1988). *Sarkastodon* from the late Eocene of Mongolia is another bearlike creodont for which a reconstruction is available (by Graham Allen for Dixon and others, 1992).

Diceratherium annectens

Only molar fragments of the rhino *Diceratherium annectens* Marsh (1873) have been found in the late Oligocene (early Arikareean) Turtle Cove Member of the John Day Formation on Carroll Rim and the footslopes of Sutton Mountain, but excellent skulls and postcranial remains continue to be found at the same stratigraphic level in and around the Sheep Rock Unit of the John Day Fossil Beds near Dayville (Peterson, 1920). Our reconstruction (Figure 10A) is based on a hornless reconstruction by Bruce Horsfall (for W.B. Scott, 1913). This Oregon rhinoceros should not be confused with a Nebraskan Miocene rhino, formerly regarded as congeneric but now assigned to *Meno-ceras cooki* (Prothero and others, 1989) and reconstructed by Bruce Horsfall (for W.B. Scott, 1913). *Diceratherium annectens* had hypsodont teeth and was one of the earliest grazing mammals (Webb, 1977).

Miohippus quartus

The three-toed horse *Miohippus quartus* Osborn (1918) is represented only by fragmentary teeth in the late Oligocene (Arikareean) Turtle Cove Member of the John Day Formation in the Painted Hills area but by fine skulls in the same stratigraphic horizon to the east near Dayville. Other species of *Miohippus* represented by skulls and complete skeletons are known from the Oligocene (Orellan and Whitneyan) Brule Formation of South Dakota (Prothero and Shubin, 1989). These and a reconstruction by Charles Halgren and associates (for Simpson and Beck, 1963) were used to prepare our version (Figure 10B). *Miohippus* like the co-occurring *Mesohippus* was a three-toed horse with the low-crowned teeth of a browser. Nevertheless, both taxa, and *Miohippus* more than *Mesohippus*, have more complex, wider, and higher crowned teeth and more elon-

gate limbs than geologically older forest horses such as *Haplohippus*, *Epihippus*, and *Orohippus*. These indicate adaptation to more open country. The distribution of fossil horses in paleosols of Oligocene age in South Dakota has suggested to MacFadden (1992), that *Mesohippus* preferred streamside woodlands, whereas *Miohippus* lived in more

open, grassy vegetation of interfluves, an idea supported to a limited extent by our discovery of *Miohippus* in paleosols in Oregon (Bestland and Retallack, 1994b).

"Entelodon" calkinsi

The large, hoglike "*Entelodon*" *calkinsi* (Sinclair) Trox-




Figure 10. Reconstructions of fossil mammals known from the late Oligocene lower Turtle Cove Member of the John Day Formation in the Painted Hills area (A and scale by R.B. Horsfall for W.B. Scott, 1913; D from Dixon and others, 1992; others original with inspiration from Sinclair, 1905; Merriam, 1906; Osborn, 1918; Peterson, 1920; W.B. Scott, 1940; Simpson and Beck, 1963; Czerkas and Glut, 1983).

ell (1920) is represented by very fragmentary remains in the late Oligocene (Arikareean) Turtle Cove Member of the John Day Formation in the Painted Hills area but by a large and complete skull at comparable stratigraphic level in the Dayville area. *Entelodon* and allied genera belong to an extinct group (Entelodontidae), distinct from but allied to pigs (Carroll, 1988). Other species of entelodons have been widely reconstructed (W.B. Scott, 1913; Czerkas and Glut, 1982; Savage and Long, 1986; Dixon and others, 1992), but our version (Figure 10F) emphasizes the distinctive pattern of bony knobs on the jaw and cheeks of the skull described by Sinclair (1905). Entelodons are unique in many ways but were probably omnivore-scavengers (Joeckel, 1990).

Hypertragulus hesperius

The small mouse deer *Hypertragulus hesperius* Hay (1902) is one of the most abundant fossil mammals in the Turtle Cove Member of the John Day Formation but is represented mainly by fragmentary remains. Good skeletons of other species of *Hypertragulus* and the similar *Leptomeryx evansi* are known from the Brule Formation of South Dakota (W.B. Scott, 1940). These and the reconstructions of *L. evansi* by Charles Knight (for Osborn, 1910) have influenced our reconstruction of *Hypertragulus hesperius* (Figure 10D). *Hypertragulus* and *Leptomeryx* are placed in separate families (Hypertragulidae and Leptomerycidae) of extinct primitive ruminants allied to the living mouse deer (Tragulidae) of Southeast Asia and Africa. Like living mouse deer, the extinct ones were probably forest browsers.

Eporeodon occidentalis

Good skulls, including the holotype of *Eporeodon occidentalis* Marsh (1873), have been found in the late Oligocene (Arikareean) Turtle Cove Member of the John Day Formation in the Painted Hills area (Thorpe, 1921). The taxonomy of these abundant Oligocene mammals remains confused (Prothero and others, 1996). *Eporeodon occidentalis* was similar to though slightly more gracile and shorter limbed than the common *Merycoidodon culbertsoni* of the Badlands of South Dakota, which has been reconstructed by Bruce Horsfall (for W.B. Scott, 1913), Charles Knight (for Osborn, 1910), and Andrew Robinson (for Dixon and others, 1992). Both the local skeletons and these reconstructions influenced our reconstruction (Figure 10C). Oreodonts are a completely extinct group of endemic North American early artiodactyls (Merycoidontidae or Oreodontidae of Carroll, 1988). They have a sheeplike appearance but are unrelated to these domestic bovids. They have also been called ruminant hogs, but pigs were a separate lineage. And although part of the larger ruminant clade, living ruminants all evolved after oreodonts. Oreodonts have a limb structure that includes feet more like paws than hooves and teeth that are low crowned. They were probably browsing animals (Bakker, 1983).

Nimravus brachyops

Only a talonid allied to these early sabre-tooth cats (Nimravidae) has been found in the late Oligocene Turtle Cove Member of the John Day Formation in the Painted Hills area, but good skulls and limb bones of *Nimravus brachyops* (Cope) Cope (1879) are known from this stratigraphic level near Dayville to the west (Toohey, 1959). The reconstruction of Graham Allen (for Dixon and others, 1992) inspired our version (Figure 10E). Nimravidae were among the earliest of the true carnivores that began to replace archaic creodont carnivores in the late Eocene. The sabre-tooth specialization of these carnivores developed independently in several different lineages of cats, the last of which became extinct at the end of the Pleistocene. *Nimravus brachyops* did not have such exaggerated canines as many later sabre-tooth cats. Its limbs were elongate and toes reduced, more like a pursuit than ambush predator, though not so cursorial as the living cheetah (Merriam, 1906).

CONCLUSION

The John Day Fossil Beds National Monument conserves a fossil record of life and landscapes in central Oregon that continues to intrigue amateur and professional paleontologists. The quality of preservation and diversity of fossil plants and animals enables an unusually detailed view of Eocene and Oligocene life on land. Associated fossil soils allow paleoecological interpretations of the fossil assemblages, and abundant volcanic rocks allow their precise radiometric dating. With our reconstructions and supporting notes, we have attempted to introduce a few of the common extinct fossil plants and animals of central Oregon. In distilling 130 years of scientific study of these fossils we are keenly aware that much more remains to be done. John Day fossils will continue to amaze and inspire for years to come.

ACKNOWLEDGMENTS

Work was funded by contract CX-9000-1-0009 from the USDI National Park Service. We thank C. Swisher, J. Dilles, E. Taylor, M. Woodburne, and S. Manchester for useful discussion and advice. Art work is by author Retallack, who takes sole responsibility for its shortcomings.

REFERENCES CITED

- Ashwill, M., 1983, Seven fossil floras in the rain shadow of the Cascade Mountains, Oregon: *Oregon Geology*, v. 45, no. 10, p. 107-111.
- Bakker, R.T., 1983, The deer flees, the wolf pursues: Incongruencies in predator-prey coevolution, in Futuyma, D.J., and Slatkin, M., eds., *Coevolution*: Sunderland, Sinauer, p. 350-382.
- Bartholomew, B., Boufford, D.E., and Spongberg, S.A., 1983, *Metasequoia ghyostroboides*—its present status in central China: *Journal of the Arnold Arboretum*, v. 64, p. 105-128.
- Basinger, J.F., 1981, The vegetative body of *Metasequoia milleri* from the middle Eocene of British Columbia: *Canadian Journal of Botany*, v. 59, p. 2379-2410.
- , 1984, Seed cones of *Metasequoia milleri* from the middle Eocene of southern British Columbia: *Canadian Journal of Botany*, v. 62, p. 281-289.
- Berry, E.W., 1928, *Stones of Celtis* in the Tertiary of the western United States: *American Museum Novitates*, v. 298, 5 p.
- Best, G.R., Hamilton, D.B., and Auble, G.J., 1984, An old growth cypress stand in Okefenokee Swamp, in Cohen, A.D., Casagrande, D.J., Andryko, M.J., and Best, G.R., eds., *The Okefenokee Swamp*. Los Alamos, N. Mex., Wetlands Surveys, p. 132-143.
- Bestland, E.A., and Retallack, G.J., 1994a, Geology and paleoenvironments of the Clamo Unit, John Day Fossil Beds National Monument, Oregon: Unpublished final report to the USDI National Park Service for Contract CX-9000-1-0009, 160 p.
- , 1994b, Geology and paleoenvironments of the Painted Hills Unit, John Day Fossil Beds National Monument, Oregon: Unpublished final report to the USDI National Park Service for Con-

- tract CX-9000-1-0009, 211 p.
- Bestland, E.A., Retallack, G.J., and Fremd, T., 1994, Sequence stratigraphy of the Eocene-Oligocene transition: Examples from the nonmarine, volcanically influenced John Day Basin, in Swanson, D.A., and Haugerud, R.A., eds., *Geologic field trips in the Pacific Northwest*: Seattle, Wash., University of Washington, v. 1, p. A1-A19.
- 1995, Geology of the late Eocene Clarno Unit, John Day Fossil Beds National Monument, central Oregon, in Santucci, V.L., and McClelland, L., eds., *National Park paleontological research: USDI National Park Service Technical Report NPS/NRPO/NRTR 95/16*, p. 66-72.
- Bestland, E.A., Retallack, G.J., Rice, A.E., and Mindszenty, A., 1996, Late Eocene detrital laterites in central Oregon: Mass balance geochemistry, depositional setting, and landscape evolution: *Geological Society of America Bulletin*, v. 108, no. 3, p. 285-302.
- Bones, T.J., 1979, Atlas of fossil fruits and seeds from north-central Oregon: Portland, Oreg., Oregon Museum of Science and Industry Occasional Paper 1, 23 p.
- Breedlove, D.E., 1973, The phytogeography and vegetation of Chiapas (Mexico), in Graham, A.E., ed., *Vegetation and vegetational history of northern Latin America*: Amsterdam, Elsevier, p. 149-165.
- Brown, J.T., 1975, *Equisetum clarnoi*, a new species based on petrifications from the Eocene of Oregon: *American Journal of Botany*, v. 62, p. 410-415.
- Burger, W.C., 1983, *Alnus acuminata* (jaul, alder), in Janzen, D.H., ed., *Costa Rican natural history*: Chicago, Ill., University of Chicago Press, p. 188-189.
- Burnham, R.J., and Spicer, R.A., 1986, Forest litter preserved by volcanic activity at El Chichón, Mexico: A potentially accurate record of the pre-eruption vegetation: *Palaïos*, v. 1, p. 158-161.
- Carroll, R.L., 1988, Vertebrate paleontology and evolution: New York, Freeman, 698 p.
- Cevallos-Ferriz, S.R.S., and Stockey, R.A., 1988a, Permineralized fruits and seeds from the Princeton chert (middle Eocene) of British Columbia: *Lythraceae*: *Canadian Journal of Botany*, v. 63, p. 303-312.
- 1988b, Permineralized fruits and seeds from the Princeton chert (middle Eocene) of British Columbia: *Araceae*: *American Journal of Botany*, v. 75, p. 1099-1113.
- 1989, Permineralized fruits and seeds from the Princeton chert (middle Eocene) of British Columbia: *Nymphaeaceae*: *Botanical Gazette*, v. 150, p. 207-217.
- 1990a, Permineralized fruits and seeds from the Princeton chert (middle Eocene) of British Columbia: *Vitaceae*: *Canadian Journal of Botany*, v. 68, p. 288-295.
- 1990b, Vegetative remains of the Magnoliaceae from the Princeton chert (middle Eocene) of British Columbia: *Canadian Journal of Botany*, v. 68, p. 1327-1339.
- 1991, Permineralized fruits and seeds from the Princeton chert (middle Eocene) of British Columbia: *Prunioideae*: *Botanical Gazette*, v. 152, p. 369-379.
- Cevallos-Ferriz, S.R.S., Erwin, D.M., and Stockey, R.A., 1993, Further observations on *Palaeoecora similkameensis* (Rosaceae) from the middle Eocene Princeton chert of British Columbia, Canada: *Review of Palaeobotany and Palynology*, v. 78, p. 277-291.
- Chaney, R.W., 1924, Quantitative studies of the Bridge Creek flora: *American Journal of Science*, v. 8, p. 127-144.
- 1925a, A comparative study of the Bridge Creek Flora and the modern redwood forest: *Carnegie Institute of Washington Publication* 349, p. 3-22.
- 1925b, Notes on two fossil hackberries from the Tertiary of the western United States: *Carnegie Institute of Washington Publication* 349, p. 51-56.
- 1937, Cycads from the upper Eocene of Oregon [abs.]: *Geological Society of America Proceedings for 1936*, p. 397.
- 1938, Paleoecological interpretations of Cenozoic plants in western North America: *Botanical Review*, v. 4, p. 371-396.
- 1948a, The ancient forests of Oregon: Eugene, Oreg., Oregon State System of Higher Education Condon Lectures, 56 p.
- 1948b, The bearing of living *Metasequoia* on problems of Tertiary paleobotany: *National Academy of Science Proceedings*, v. 34, no. 11, p. 503-515.
- 1951, A revision of fossil *Sequoia* and *Taxodium* in western North America based on the recent discovery of *Metasequoia*: *American Philosophical Society Transactions*, v. 40, p. 171-263.
- 1952, Conifer dominants in the middle Tertiary of the John Day Basin, Oregon: *Palaeobotanist*, v. 1, p. 105-113.
- Chu, K., and Cooper, W.S., 1950, An ecological reconnaissance in the native home of *Metasequoia glyptostroboides*: *Ecology*, v. 31, p. 260-278.
- Clark, R.D., 1989, The odyssey of Thomas Condon: Portland, Oreg., Oregon Historical Society Press, 567 p.
- Cope, E.D., 1873, On the extinct Vertebrata of the Eocene of Wyoming, observed by the expedition of 1872, with notes on the geology: U.S. Geological and Geographical Survey of the Territories (Hayden), *Annual Report*, v. 6, p. 543-649.
- 1879, On the genera of Felidae and Canidae: *Academy of Natural Sciences of Philadelphia, Proceedings*, v. 31, p. 168-194.
- Crane, P.R., and Stockey, R.A., 1985, Growth and reproductive biology of *Joffrea speirsii* gen. et sp. nov., a *Cercidiphyllum*-like plant from the late Paleocene of Alberta, Canada: *Canadian Journal of Botany*, v. 63, p. 340-364.
- Czerkas, S.M., and Glut, D.F., 1982, Dinosaurs, mammoths and cavemen: The art of Charles R. Knight: New York, Dutton, 120 p.
- DiMichele, W.A., Phillips, T.L., and Olmstead, R.G., 1987, Opportunistic evolution: Abiotic environmental stress and the fossil record of plants: *Review of Palaeobotany and Palynology*, v. 50, p. 151-178.
- Dixon, D., Cox, B., Savage, R.J.G., and Gardner, B., 1992, The Macmillan illustrated encyclopedia of dinosaurs and prehistoric animals: New York, Macmillan, 312 p.
- Dott, R.H., and Batten, R.L., 1981, *Evolution of the Earth*: New York, McGraw-Hill, 574 p.
- Eaton, J.G., 1985, Paleontology and correlation of the Eocene Tepee Trail and Wiggins Formations in the North Fork of Owl Creek area, southeastern Absaroka Range, Hot Springs County, Wyoming: *Journal of Vertebrate Paleontology*, v. 5, p. 345-370.
- Erwin, D.M., and Stockey, R.A., 1989a, Permineralized monocotyledons from the middle Eocene Princeton chert (Allenby Formation) of British Columbia: *Alismataceae*: *Canadian Journal of Botany*, v. 67, p. 2636-2645.
- 1989b, Sapindaceous flowers from the middle Eocene Princeton chert (Allenby Formation) of British Columbia, Canada: *Canadian Journal of Botany*, v. 68, p. 2025-2034.
- 1994, Permineralized monocotyledons from the middle Eocene Princeton chert (Allenby Formation) of British Columbia: *Araceae*: *Palaeontographica*, v. B234, p. 19-40.
- Evanoff, E., Robinson, P., Murphy, P.C., Kron, D.C., Engard, D., and Monaco, P., 1994, An early Uintan fauna from Bridger E: *Journal of Vertebrate Paleontology, Supplement*, v. 14, p. 24A.
- Evernden, J.F., and James, G.T., 1964, Potassium-argon dates of the Tertiary floras of North America: *American Journal of Science*, v. 262, p. 945-974.
- Fremd, T., 1988, Assemblages of fossil vertebrates in pre-ignimbrite deposits of the Turtle Cove Member, John Day Formation (Arikarean), from outcrops within the Sheep Rock Unit, John Day Fossil Beds National Monument: *Journal of Vertebrate Paleontology, Suppl.*, v. 8, p. A15.
- 1991, Early Miocene mammalian populations from Turtle Cove, Oregon: *Journal of Vertebrate Paleontology, Supplement*, v. 11, p. A29.
- 1993, Refinement of spatial and temporal distributions of biotas within the volcaniclastic sequences of the John Day Basin: *Journal of Vertebrate Paleontology, Supplement*, v. 13, p. A36.
- Fremd, T., and Wang, X.-M., 1995, Resolving blurred faunas: Biostratigraphy in John Day Fossil Beds National Monument, in Santucci, V.L., and McClelland, L., eds., *National Park paleontological research: USDI National Park Service Technical Report NPS/NRPO/NRTR 95/16*, p. 73-79.
- Gentry, A.H., 1980, Family 109: Sabiaceae, in Woodson, R.E., and Schery, R.W., eds., *Flora of Panama: Annals of the Missouri Botanical Garden*, v. 67, p. 949-963.
- Getahun, A., and Retallack, G.J., 1991, Early Oligocene paleoenvironment of a paleosol from the lower part of the John Day Formation near Clarno, Oregon: *Oregon Geology*, v. 53, no. 6, p. 131-136.
- Gómez-Pompa, A., 1973, Ecology of the vegetation of Veracruz, in Graham, A.E., ed., *Vegetation and vegetational history of northern Latin America*: Amsterdam, Elsevier, p. 73-148.
- Gregory, I., 1969, Fossilized palm wood in Oregon: *Oregon Department of Geology and Mineral Industries, Ore Bin*, v. 31, no. 5, p. 93-110.
- Hanson, C.B., 1973, Geology and vertebrate faunas in the type area of the Clarno Formation, Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 5, no. 1, p. 50.
- 1989, *Telaceras radinskyi*, a new primitive thinoceeroid from the late Eocene Clarno Formation of Oregon, in Prothero, D.R., and Schoch, R.M., eds., *The evolution of perissodactyls*: New York, Oxford University Press, p. 235-256.
- Hartshorn, G.S., 1983, *Plants, in Janzen, D.H., ed., Costa Rican natural history*: Chicago, University of Chicago Press, p. 118-158.
- Hay, O.P., 1902, Bibliography and catalogue of the fossil Vertebrata of North America: U.S. Geological Survey Bulletin 179, 868 p.
- 1908, The fossil turtles of North America: *Carnegie Institution of Washington Publication* 75, 568 p.
- Hay, R.L., 1963, Stratigraphy and zoolitic diagenesis of the John Day Formation of Oregon: *University of California Publications in Geological Sciences*, v. 42, no. 5, p. 199-261.
- Hu, S.-Y., 1980, The *Metasequoia* flora and its phytogeographical significance: *Journal of the Arnold Arboretum*, v. 61, p. 41-94.
- Hutchinson, J.H., 1992, Western North American reptile and amphibian record across the Eocene/Oligocene boundary and its climatic implications, in Prothero, D.R., and Berggren, W.A., eds., *Eocene-Oligocene climatic and biotic evolution*: Princeton, N.J., Princeton University Press, p. 451-463.
- Joekel, R.M., 1990, A functional interpretation of the masticatory system and paleoecology of entelodonts: *Paleobiology*, v. 16, p. 459-482.
- Kitts, D.B., 1957, A revision of the genus *Orohippus* (Perissodactyla, Equidae): *American Museum Novitates*, v. 1864, p. 1-40.
- Khucking, E.P., 1956, The fossil Betulaceae of western North America: Berkeley, Calif., University of California master's thesis, 169 p.
- Knowlton, F.H., 1902, Fossil flora of the John Day Basin, Oregon: U.S. Geological Survey Bulletin 204, 153 p.
- Kvacek, Z., Buzek, C., and Manchester, S.R., 1991, Fossil fruits of *Pteleacarpum* Weyland—*Tiliaceae* not *Sapindaceae*: *Botanical Gazette*, v. 15, p. 522-523.
- Lander, E.B., 1977, A review of the Oreodonta (Mammalia, Artiodactyla): Berkeley, Calif., University of California doctoral dissertation, 472 p.
- Leidy, J., 1871, Remarks on fossil vertebrates from Wyoming: *Academy of Natural Sciences of Philadelphia, Proceedings*, v. 22, p. 228-229.
- Leopold, E.B., Liu, G.-W., and Clay-Poole, S., 1992, Low biomass vegetation in the Oligocene?, in Prothero, D.R., and Berggren, W.A., eds., *Eocene-Oligocene climatic and biotic evolution*: Princeton, N.J., Princeton University Press, p. 399-420.
- Lucas, S.G., 1992, Redefinition of the Duchesnean Land Mammal "Age," late Eocene of western North America, in Prothero, D.R., and Berggren, W.A., eds., *Eocene-Oligocene climatic and biotic evolution*: Princeton, N.J., Princeton University Press, p. 88-105.
- MacFadden, B.J., 1992, Fossil horses: Systematics, paleobiology, and evolution of the family Equidae: New York, Cambridge University Press, 369 p.
- MacGinitie, H.D., 1969, The Eocene Green River flora of northwestern Colorado and northeastern Utah: *University of California Publications in Geological Sciences*, v. 83, 203 p.
- Mader, B., 1989, The Brontotheriidae: A systematic revision and preliminary phylogeny of North American genera, in Prothero, D.R., and Schoch, R.M., eds., *The evolution of Perissodactyls*: New York, Oxford University Press, p. 458-484.
- Manchester, S.R., 1977, Wood of *Tapirtria* (Anacardiaceae) from the Paleogene Clarno Formation of Oregon: *Review of Palaeobotany and Palynology*, v. 23, p. 119-127.
- 1979, *Triplochitoxylon* (Sterculiaceae): A new genus of wood from the Eocene of Oregon and its bearing on the xylem evolution of the extant genus *Triplochiton*: *American Journal of Botany*, v. 66, p. 699-708.
- 1980a, *Chattawayia* (Sterculiaceae): A new genus of wood from the Eocene of Oregon and its implications for xylem evolution of the extant genus *Pterospermum*: *American Journal of Botany*, v. 67, p. 58-67.
- 1980b, Leaves, wood, and fruits of *Meliosma* from the Eocene Clarno Formation of Oregon [abs.]: *Botanical Society of America Miscellaneous Series Publication* 158, p. 70.
- 1981, Fossil plants of the Eocene Clarno nut beds: *Oregon Geology*, v. 43, no. 6, p. 75-81.
- 1983, Fossil wood of the Engelhardiaceae (Juglandaceae) from the Eocene of North America: *Engelhardioxylon* gen. nov.: *Botanical Gazette*, v. 144, p. 157-163.
- 1986, Vegetative and reproductive morphology of an extinct plane tree (Platanaceae) from the Eocene of western North America: *Botanical Gazette*, v. 147, p. 200-226.
- 1987, The fossil history of the Juglandaceae: *Monographs from the Missouri Botanical Garden*, v. 21, 137 p.
- 1988, Fruits and seeds of *Tapiscia* (Staphyleaceae) from the middle Eocene of Oregon, U.S.A.: *Tertiary Research*, v. 9, p. 59-66.
- 1991, *Cruciptera*, a new Juglandaceous winged fruit from the Eocene and Oligocene of western North America: *Systematic Botany*, v. 16, p. 715-723.
- 1992, Flowers, fruits, and pollen of *Florisantia*, an extinct malvacean genus from the Eocene and Oligocene of western North America: *American Journal of Botany*, v. 79, p. 996-1008.
- 1994a, Fruits and seeds of the middle Eocene Nut Beds Flora, Clarno Formation, Oregon: Ithaca, N.Y., Paleontological Research Institution, *Palaeontographica Americana* 58, 205 p.
- 1994b, Inflorescence bracts of fossil and extant *Tilia* in North America, Europe, and Asia: Patterns of morphologic divergence and biogeographic history: *American Journal of Botany*, v. 81, p. 1176-1185.
- 1995, Yes, we had bananas: *Oregon Geology*, v. 57, no. 2, p. 41-43.
- Manchester, S.R., Collinson, M.E., and Goth, K., 1994, Fruits of the Juglandaceae from the Eocene of Messel, Germany, and implications for early Tertiary phytogeographic exchange between Europe and western North America: *International Journal of Plant Science*, v. 155, p. 388-394.
- Manchester, S.R., and Kress, W.J., 1993, Fossil bananas (Musaceae): *Ensete oregonense* sp. nov. from the Eocene of western North America and its phytogeographic significance: *American*

- Journal of Botany, v. 80, p. 1264-1272.
- Manchester, S.R., and Meyer, H.W., 1987, Oligocene plants of the John Day Formation, Fossil, Oregon. *Oregon Geology*, v. 49, p. 115-127.
- Manchester, S.R., and Wheeler, E.A., 1993, Extinct Juglandaceae wood from the Eocene of Oregon and its implications for xylem evolution in the Juglandaceae: *Journal of the International Association of Wood Anatomists*, v. 14, p. 103-111.
- Marsh, O.C., 1873, Notice of new Tertiary mammals: *American Journal of Science*, v. 5, p. 407-410, 485-488.
- 1874, Notice of new equine mammals from the Tertiary formation: *American Journal of Science*, v. 6, p. 247-258.
- 1875, Ancient lake basins of the Rocky Mountain region: *American Journal of Science*, v. 9, p. 45-52.
- 1878, Vertebrates of the Bridger, in King, C., *Systematic geology: U.S. geological exploration of the fortieth parallel*, annual report, p. 403-405.
- McGrew, P.O., 1953, A new and primitive early Oligocene horse from trans-Pecos Texas: *Fieldiana Geology*, v. 10, p. 167-171.
- McKee, T.M., 1970, Preliminary report on fossil fruits and seeds from the Mammal Quarry of the Clarno Formation, Oregon: *Oregon Department of Geology and Mineral Industries, Ore Bin*, v. 32, no. 7, p. 117-132.
- Mellet, J.S., 1969, A skull of *Hemipalodon* (Mammalia, order Deltatheridia) from the Clarno Formation of Oregon: *American Museum Novitates*, v. 2387, 19 p.
- Merriam, J.C., 1901a, A geological section through the John Day Basin: *Journal of Geology*, v. 9, p. 71-72.
- 1901b, A contribution to the geology of the John Day Basin: *University of California Publications in Geological Sciences*, v. 2, p. 269-314.
- 1906, Carnivora from the Tertiary formation of the John Day region: *University of California Publications in Geological Sciences*, v. 5, p. 1-64.
- Merriam, J.C., and Sinclair, W.J., 1907, Tertiary faunas of the John Day region: *University of California Publications in Geological Sciences*, v. 5, p. 171-205.
- Merrill, E.D., 1948, *Metasequoia* another "living fossil": *Arnoldia*, v. 8, p. 1-6.
- Miki, S., 1941, On the change of flora in eastern Asia since Tertiary Period (I). The clay or lignite beds flora in Japan, with special reference to the *Pinus trifolia* beds in central Hondo: *Japanese Journal of Botany*, v. 11, p. 260-262.
- Newberry, J.S., 1883, Brief descriptions of fossil plants, chiefly Tertiary, from western North America: *U.S. National Museum, Proceedings*, v. 5, p. 502-514.
- 1898, The later extinct floras of North America: *U.S. Geological Survey Monograph 35*, 295 p.
- Osborn, H.F., 1910, *The age of mammals in Europe, Asia, and North America*: New York, Macmillan, 635 p.
- 1918, Equidae of the Oligocene, Miocene, and Pliocene of North America: *Iconographic type revision: American Museum of Natural History Memoir 2*, 326 p.
- 1929, The titanotheres of ancient Wyoming, Dakota, and Nebraska: *U.S. Geological Survey Monograph 55*, 953 p.
- Owen-Smith, R.N., 1988, *Megaherbivores*: New York, Cambridge University Press, 369 p.
- Peattie, D.C., 1991, *A natural history of trees of eastern and central North America*: Boston, Mass., Houghton Mifflin, 751 p.
- Peterson, O.A., 1920, The American diceratheres: *Carnegie Museum Memoir 7*, p. 399-488.
- Porter, D.M., 1973, The vegetation of Panama—a review, in Graham, A., ed., *Vegetation and vegetational history of northern Latin America*: Amsterdam, Elsevier, p. 167-201.
- Pratt, J.A., 1988, Paleoenvironment of the Eocene/Oligocene Hancock Mammal Quarry, upper Clarno Formation, Oregon: Eugene, Ore., University of Oregon master's thesis, 104 p.
- Prothero, D.R., Guérin, C., and Manning, E., 1989, The history of the Rhinocerotidae, in Prothero, D.R., and Schoch, R.M., eds., *The evolution of perissodactyls*: New York, Oxford University Press, p. 321-340.
- Prothero, D.R., Heaton, T.H., and Stanley, S.M., 1996, Speciation and faunal stability in mammals during the Eocene-Oligocene climatic crash: *Paleogeography, Paleoclimatology, Paleoeecology* (in press).
- Prothero, D.R., and Shubin, N., 1989, The evolution of Oligocene horses, in Prothero, D.R., and Schoch, R.M., eds., *The evolution of perissodactyls*: New York, Oxford University Press, p. 142-175.
- Prothero, D.R., and Swisher, C.C., 1992, Magnetostratigraphy and geochronology of the terrestrial Eocene-Oligocene transition in North America, in Prothero, D.R., and Berggren, W.A., eds., *Eocene-Oligocene climatic and biotic evolution*: Princeton, N.J., Princeton University Press, p. 46-73.
- Radinsky, L.B., 1963, Origin and evolution of North American Tapiroidea: *Peabody Museum of Natural History Bulletin*, v. 17, p. 1-106.
- Retallack, G.J., 1981, Preliminary observations on fossil soils in the Clarno Formation (Eocene to early Oligocene) near Clarno, Oregon: *Oregon Geology*, v. 43, no. 11, p. 147-50.
- 1991a, A field guide to mid-Tertiary paleosols and paleoclimatic changes in the high desert of central Oregon—Part 1: *Oregon Geology*, v. 53, no. 3, p. 51-59.
- 1991b, A field guide to mid-Tertiary paleosols and paleoclimatic changes in the high desert of central Oregon—Part 2: *Oregon Geology*, v. 53, no. 4, p. 75-80.
- 1992, Paleosols and changes in climate and vegetation across the Eocene-Oligocene boundary, in Prothero, D.R., and Berggren, W.A., eds., *Eocene-Oligocene climatic and biotic evolution*: Princeton, N.J., Princeton University Press, p. 383-398.
- 1994a, The environmental factor approach to the interpretation of paleosols, in Amundson, R., Harden, J., and Singer, M., eds., *Factors of soil formation: A fiftieth anniversary perspective*: Soil Sciences Society of America, Special Publications, v. 33, p. 31-64.
- 1994b, A pedotype approach to latest Cretaceous and earliest Tertiary paleosols in eastern Montana: *Geological Society of America Bulletin*, v. 106, no. 11, p. 1377-1397.
- Retallack, G.J., and Dilcher, D.L., 1981, A coastal hypothesis for the dispersal and rise to dominance of flowering plants, in Niklas, K.J., ed., *Paleobotany, paleoecology, and evolution*: New York, Praeger, v. 2, p. 27-77.
- Rice, A.E., 1994, A study of the late Eocene paleoenvironment in central Oregon based on paleosols from the lower Big Basin Member of the John Day Formation in the Painted Hills: Eugene, Ore., University of Oregon master's thesis, 154 p.
- Robinson, P.T., Brem, G.F., and McKee, E.H., 1984, John Day Formation of Oregon: A distal record of early Cascade volcanism: *Geology*, v. 12, no. 4, p. 229-232.
- Rothwell, G.W., and Basinger, J.F., 1979, *Metasequoia milleri* n. sp., anatomically preserved pollen cones from the middle Eocene (Allenby Formation) of British Columbia: *Canadian Journal of Botany*, v. 57, p. 958-970.
- Russell, L.L., 1938, The skull of *Hemipalodon grandis*, a giant Oligocene creodont: *Royal Society of Canada Transactions*, v. 32, p. 61-66.
- Savage, R.J.G., and Long, M.R., 1986, *Mammal evolution*: New York, Facts on File, 259 p.
- Scheele, W.E., 1955, *The first mammals*: New York, World, 128 p.
- Schoch, R.M., 1989, A review of tapiroids, in Prothero, D.R., and Schoch, R.M., eds., *The evolution of perissodactyls*: New York, Oxford University Press, p. 298-320.
- Schopf, J.M., 1948, Should there be a living *Metasequoia*? *Science*, v. 107, p. 344-345.
- Scott, R.A., 1954, Fossil fruits and seeds from the Eocene Clarno Formation of Oregon: *Paleontographica*, v. B96, p. 66-97.
- 1955, *Cryptocolax*—a new genus of fungi (Aspergillaceae) from the Eocene of Oregon: *American Journal of Botany*, v. 43, p. 589-593.
- 1956, Evolution of some endocarpal features in the tribe Tinosporeae (Menispermaceae): *Evolution*, v. 10, p. 74-81.
- Scott, R.A., and Barghoorn, E.S., 1956, The occurrence of *Euptelea* in the Cenozoic of western North America: *Journal of the Arnold Arboretum*, v. 36, p. 259-266.
- Scott, R.A., Barghoorn, E.S., and Prakash, U., 1962, Wood of *Ginkgo* in the Tertiary of western North America: *American Journal of Botany*, v. 49, p. 1095-1101.
- Scott, R.A., and Wheeler, E.A., 1982, Fossil woods from the Eocene Clarno Formation of Oregon: *International Association of Wood Anatomists (IAWA) Bulletin*, v. 3, p. 135-154.
- Scott, W.B., 1913, *A history of land mammals in the western hemisphere*: New York, Macmillan, 693 p.
- 1940, The mammalian fauna of the White River Oligocene. Part IV. Artiodactyla: *American Philosophical Society Transactions*, v. 28, p. 159-172.
- 1941, The mammalian fauna of the White River Oligocene. Part V. Perissodactyla: *American Philosophical Society Transactions*, v. 28, p. 747-980.
- 1945, The mammalia of the Duchesne River Oligocene: *American Philosophical Society Transactions*, v. 34, p. 209-253.
- Simpson, G.G., and Beck, W.S., 1963, *Life: an introduction to biology*: New York, Harcourt, Brace and World, 869 p.
- Sinclair, W.J., 1905, New or imperfectly known rodents and ungulates from the John Day series: *University of California Publications, Bulletin of the Department of Geological Sciences*, v. 47, p. 587-590.
- Smith, G.S., 1988, Paleoenvironmental reconstruction of Eocene fossil soils from the Clarno Formation in eastern Oregon: Eugene, Ore., University of Oregon master's thesis, 167 p.
- Spinar, Z.V., and Burian, Z., 1972, *Life before Man*: London, Thames and Hudson, 228 p.
- Stirton, R.A., 1944, A rhinoceros tooth from the Clarno Eocene of Oregon: *Journal of Paleontology*, v. 18, p. 265-267.
- Stock, C., 1936, Titanotheres from the Titus Canyon Formation, California: *National Academy of Sciences, Proceedings*, v. 22, p. 656-661.
- Stockey, R.A., 1987, A permineralized flower from the middle Eocene of British Columbia, Canada: *American Journal of Botany*, v. 74, p. 1878-1887.
- Stockey, R.A., and Pigg, K.B., 1991, Flowers and fruits of *Princetonia allenbyensis* (Magnoliopsida, family indet.) from the middle Eocene Princeton chert of British Columbia: *Reviews of Paleobotany and Palynology*, v. 70, p. 163-172.
- Swisher, C.C., and Prothero, D.R., 1990, Single crystal $^{40}\text{Ar}/^{39}\text{Ar}$ dating of the Eocene-Oligocene transition in North America: *Science*, v. 249, p. 760-762.
- Tanai, T., and Wolfe, J.A., 1977, Revisions of *Ulmus* and *Zelkova* in the middle and late Tertiary of western North America: *U.S. Geological Survey Professional Paper 1026*, 14 p.
- Thorpe, M.R., 1921, The John Day eoprodons, with descriptions of new genera and species: *American Journal of Science*, v. 2, p. 93-111.
- Toohy, L., 1959, The species of *Nimravus* (Carnivora, Felidae): *American Museum of Natural History Bulletin*, v. 118, p. 75-112.
- Troxell, E.L., 1920, Entelodonts in the Marsh collection: *American Journal of Science*, v. 50, p. 243-255.
- van Beusekom, C.F., and van de Water, T.P.M., 1989, Sabiaceae: *Flora malesiana*, series 1, v. 10, no. 4, p. 679-715.
- Vance, J.A., 1988, New fission-track and K-Ar ages from the Clarno Formation, Challis age volcanic rocks in north-central Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 20, no. 6, p. 473.
- Walker, G.W., 1977, Geologic map of Oregon east of the 121st meridian: *U.S. Geological Survey Miscellaneous Investigations Series Map I-902*, scale 1:500,000.
- Walker, G.W., and Robinson, P.T., 1990, Paleocene(?) Eocene, and Oligocene(?) rocks of the Blue Mountains region, *chap. 2 of Walker, G.W., ed., Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Cenozoic geology of the Blue Mountains region*: U.S. Geological Survey Professional Paper 1437, p. 13-27.
- Webb, S.D., 1977, A history of savanna vertebrates in the New World. Part 1. North America: *Annual Reviews of Ecology and Systematics*, v. 8, p. 355-380.
- Wheeler, E.R., Scott, R.A., and Barghoorn, E.S., 1977, Fossil dicotyledonous woods from Yellowstone National Park: *Journal of the Arnold Arboretum*, v. 58, p. 280-306.
- White, J.D.L., and Robinson, P.T., 1992, Intra-arc sedimentation in a low-lying marginal arc, Eocene Clarno Formation, central Oregon: *Sedimentary Geology*, v. 80, p. 89-114.
- Wolfe, J.A., 1981, Paleoclimatic significance of the Oligocene and Neogene floras of the northwestern United States, in Niklas, K.J., ed., *Paleobotany, paleoecology, and evolution*: New York, Praeger, v. 2, p. 79-101.
- 1987, An overview of the origins of the modern vegetation and flora of the northern Rocky Mountains: *Annals of the Missouri Botanical Garden*, v. 74, p. 785-803.
- 1992, Climatic, floristic, and vegetational changes over the Eocene/Oligocene boundary in North America, in Prothero, D.R., and Berggren, W.A., eds., *Eocene-Oligocene climatic and biotic evolution*: Princeton, N.J., Princeton University Press, p. 421-436.
- Wolfe, J.A., and Tanai, T., 1987, Systematics, phylogeny and distribution of *Acer* (maples) in the Cenozoic of western North America: *Journal of the Faculty of Science, Hokkaido University*, v. 22, p. 1-246.
- Wood, C.E., 1974, *A student's atlas of flowering plants: Some dicotyledons of eastern North America*: New York, Harper and Row, 120 p. □

John Day Fossil Beds National Monument has a home page on the Internet

Those interested in finding out more about the John Day Fossil Beds and ready to get on the Internet can find the home page of the John Day Fossil Beds National Monument under the address <http://www.nps.gov/joda>.

Without the final "joda" this address is for the National Park Service home page, which can lead you to many other sources of information on National Parks. □

BOOK REVIEWS

Fruits and Seeds of the Middle Eocene Nut Beds Flora, Clarno Formation, Oregon, by Steven R. Manchester. Ithaca, N.Y., Paleontological Research Institution, *Palaeontographica Americana* 58, August 24, 1994, 205 p., including 70 plates, \$65.00.

Reviewed by Ted Fremd, Paleontologist, John Day Fossil Beds National Monument, HCR 82 Box 126, Kimberly, OR 97848, e-mail Ted_Fremd@NPS.GOV

The predominantly volcanoclastic Clarno Formation, first recognized and described around the turn of the century (Merriam, 1901), covers a huge expanse of eastern Oregon. Manchester's excellent new monograph focuses on only a tiny portion of the record preserved in these strata. The Nut Beds flora is found in 44-million-year-old deposits at a minuscule site within the boundaries of what is now the Clarno Unit of John Day Fossil Beds National Monument. Ever since its discovery, this peculiar locality has puzzled and delighted many botanical workers. In 1942, a fossil enthusiast named Thomas Bones (who did not work with vertebrates) began diligently working these beds. By 1954, when the first monograph on the flora appeared (Scott, 1954), Bones had laboriously collected and painstakingly prepared microscopic specimens of fossil seeds and fruits from over 10 tons of material. Bones continued collecting for another 30 years, providing a classic example of the role "amateurs" can have in paleontology. He amassed significant collections housed today in major repositories where they are available for study, and he published a photographic atlas of his specimens (Bones, 1979).

Steve Manchester's new publication is an exhaustive and thorough summary of the Nut Bed fossil seeds and fruits. It is based on new examinations of over 20,000 specimens, most of them from Bones' collections. Manchester's effort is unique in the North American paleobotanical literature and has produced a huge success, particularly since the Nut Beds site has produced extraordinarily well preserved fruits, flowers, pollen, leaves, stems, roots, and other tissue systems in association.

A marvelously complex picture of a diverse (145 genera), paratropical flora emerges from Manchester's work. Vines or other climbing plants make up nearly half of the flora for which the growth habit can be deduced. A puzzling mix of palms, ferns, cycads, and numerous types of flowering plants is described (with only one herbaceous angiosperm, the banana *Ensete*, detailed in Manchester and Kress, 1993). For years, we have been comparing the Clarno rain forest to habitats such as those found in modern southeast Asia, but comparisons with modern forests fail to do justice to the diversity and degree of difference of the Clarno rain forest: it appears here much more bizarre than anybody had thought so far. If modern botanists could somehow walk through the Clarno jungle, they would not just see alien species—they would be unable to assign over

half of the plants to any of the modern families!

A small but equally odd suite of vertebrates is also known from the Nut Beds, including crocodylians, primitive horses, and brontotheres, some of which undoubtedly feasted on the fruits and helped to disperse the plants.

Professionals and amateurs alike will be delighted with the great number and quality of the photographic plates in Manchester's book. I counted 1,155 clear, crisp images of 170 species with excellent descriptions, including whole-rock specimens, acetate peels, stereopairs, and beautiful scanning-electron microscopy. Incidentally, these photographs would be very useful on a CD-ROM linked to the text descriptions and illustrations, as several publishers are finding. The bulk of the text (90 of 120 pages) is concerned with descriptions of the taxa, including dozens of new species and/or combinations, following standard systematic procedure. The descriptions are accompanied by discussions, clear text figures, and well-designed tables that do an excellent job of clarifying synonymies.

Among the figured specimens are some that may correspond to insect reproductive structures rather than plants. Plate 66, for example, includes illustrations of "seeds" that I had assumed were brood chambers of one or more species of burrowing hymenopteran that are also found in younger John Day Basin strata. In recognizing the possibility of incorrect identification, Manchester decided that, "Rather than exclude taxa that could possibly be construed as insect eggs, I believe it is useful to document all of the structures because of the possible importance for biostratigraphy and/or future insect work" (p. 18). Still, the consequence is somewhat odd: One may have to consult botanical descriptions of micropyles, seed coats, and endosperm cavities for what may be entomological features.

Incidentally, paleobotanists are not the first organisms tempted into thinking that some of the little shapes we find today in Clarno muds are plant seeds. Modern species of "stick insects," of which the oldest known fossil eggs are found in the Nut Beds (Sellick, 1994), have evolved a complex mimicry: They succeed in fooling some ants into thinking the stick insect eggs are a type of plant seed. The ants carry these "seeds" into their nests, unwittingly assisting the reproductive efforts of the stick insects—as presumably their Eocene counterparts did.

Not much can be found that detracts from the excellence of Manchester's work. Among some minor flaws is the erroneous location in text figure 3 of the Hancock Quarry, placed in an outcrop rather far from the actual site. The pages devoted to the geology contain several antiquated interpretations of what is admittedly a puzzling stratigraphic sequence, and they are only partially corrected by "Notes added in proof." For example, instead of the earlier identification of the Nut Beds as "predating an andesitic plug" (p. 9), it is now clear that the Nut Beds actually are high in a section of lahars that accumulated on top of the plug (which is actually dacite) (Bestland and others, 1994). Manchester dismisses the long-standing "lake delta" hy-

pothesis and rightly points out that the evidence is more consistent with a flood/slurry deposit associated with a lahar.

Unfortunately, the binding is pathetically inadequate for a book of this caliber. Since it will often serve as a reference, users will have to reinforce the binding to guard against losing individual pages.

Fruits and Seeds of the Middle Eocene Nut Beds Flora is a major contribution to paleobotany and an indispensable reference tool that belongs on the shelves of every paleobotanist and worker who deals with any aspect of the terrestrial Tertiary.

References cited


- Bestland, E.A., Retallack, G.J., and T. Fremd, 1995, Geology of the late Eocene Clarno Unit, John Day Fossil Beds National Monument, central Oregon: USDI National Park Service Technical Report NPS/NRPO/NRTR 95/16, p. 66–72.
- Bones, T.J., 1979, Atlas of fossil fruits and seeds from north-central Oregon: Occasional Papers of the Oregon Museum of Science and Industry 1, 23 p.
- Manchester, S.R., and Kress, W.J., 1993, Fossil bananas (Musaceae): *Ensete oregonense* sp. nov. from the Eocene of western North America and its phylogeographic significance: American Journal of Botany, v. 80, no. 1, p. 1264–1272.
- Merriam, J.C., 1901, A contribution to the geology of the John Day Basin: University of California Publications in Geological Sciences, v. 2, p. 269–314.
- Scott, R.A., 1954, Fossil fruits and seeds from the Eocene Clarno Formation of Oregon: Paleontographica, v. B96, p. 66–97.
- Sellick, J.T.C., 1994, Phasmida (stick insect) eggs from the Eocene of Oregon: Paleontology, v. 37, pt. 4, p. 913–921.

A Rock Grows Up, by R. and M. Goodrich. Lake Oswego, Ore., GeoQuest Publishing, 56 p., \$9.95.

A Rock Grows Up is the title of a children's book, created by Lake Oswego high school teacher Michael Goodrich and his wife Randi and delightfully illustrated by Michele Han, honor student at Lake Oswego High School. The book introduces earth science concepts in such a way that they become appealing and easily comprehensible to children at the elementary school level.

Following the story of Barry Basalt, the rock that grows up, young readers of the large-print book meet a dozen characters—family members and neighbors—that represent rocks and rock-forming minerals, such as Larry Gneiss and Annie Augite. They all participate in the great geologic cycle of rock formation and decomposition that starts in the asthenosphere below a sea-floor spreading ridge and leads through plate subduction, volcanism, weathering, sedimentation, and metamorphism—from formlessness into form and back.

The whimsical pictures of the characters in the story are accompanied by schematic block diagrams that illustrate the geologic processes as they occur in the earth from the Juan de Fuca Plate to the Cascade Range—once even to Picture Gorge. In this sense, the book addresses children of Oregon and, to a lesser extent, the Pacific Northwest, conveying the framework of basic geology these children can experience at home.



Cover of *A Rock Grows Up*

The about fifty pages of the book include also a glossary of geologic terms and an index, also a list of significant technical books that introduce the teacher to the geology of the Pacific Northwest and to the major processes touched upon in the story of Barry Basalt. A bit of caution is advised in following the pronunciation aids given throughout the text and the glossary. A teacher may want to try them out before introducing them to children of the Northwest.

A Rock Grows Up is conceived as a teaching tool and thus accompanied by more teaching aids: a curriculum guide for student activities in lab exercises, coloring, and creative writing and with more materials for the use of the teacher (\$7.95). The authors also offer a kit containing tagged samples of the rocks and minerals of the story (\$14.95).

The book and accompanying materials are available from the publishers, GeoQuest Publishing, P.O. Box 1665, Lake Oswego, OR 97035, phone (503) 635-4420; or from the Nature of the Northwest Information Center on the ground floor of the State Office Building in Portland (see address on last page of this issue). □

Offshore volcanic activity reported

The Nature of the Northwest Information Center has received numerous inquiries about volcanic activity off the Oregon coast. Internet users can find up-to-date information on this subject in the U.S. Department of Commerce NOAA/PMEL/VENTS Program under the address <http://www.pmel.noaa.gov/vents/eruption.html>. □

Oil and gas exploration and development in Oregon, 1995

by Dan E. Wermiel, Petroleum Geologist, Oregon Department of Geology and Mineral Industries

ABSTRACT

Oil and gas leasing activity was about the same during 1995 as it was in 1994. Four U.S. Bureau of Land Management (BLM) lease sales were held, and offers were received on three tracts comprising 8,250 acres located in Jefferson County, Oregon. A total of 11,760 federal acres were under lease at year's end. The State of Oregon conducted no lease auctions during the year. A total of 16 State of Oregon tracts were under lease at year's end and consisted of 25,520 acres, which is the same as 1994. Columbia County held no lease sales during 1995.

During 1995, no exploratory wells were drilled in Oregon, primarily due to the fact that the operator of Mist Gas Field, Nahama and Weagant Energy Company, had filed for bankruptcy during 1994. The field was put for sale, and Enerfin Resources was the successful bidder and will become operator of the majority of the wells at the field upon finalization of the ownership transfer. Oregon Natural Gas Development Company acquired ownership of 10 wells and certain leases at Mist Gas Field.

At the Mist Gas Field, 18 wells were productive during 1995. A total of 2.5 Btu of gas was produced with a total value of \$1.8 million during the year. This is a significant decline from the 4.2 Bcf of gas with a total value of \$6.4 million produced during 1994. Reasons for the decline in production during 1995 include the normal decline in the wells at the field; the record low of prices paid for gas, which discouraged exploration and production activities; and the decommissioning of the Nitrogen Rejection Unit during 1995, which caused several low-Btu wells to be suspended.

Final reports on the DOGAMI five-year study of the oil, gas and coal resource potential of the Tyee Basin should be published during 1996. No changes to statutes and administrative rules were made during the year.

LEASING ACTIVITY

There was little oil and gas leasing activity in Oregon during 1995. This is a continuation of a generally declining trend in leasing activity that began during the late 1980s. Activity included four public sales by the BLM at which a total of three offers were received for oil and gas leases on lands in Jefferson County. Robert F. Harrison, Seattle, Washington, applied for 8,250 acres during December 1995. He and other individuals have held leases on a block of federal land comprising about 11,680 acres in this general area since 1977. This acreage, plus an 80-acre parcel located in Columbia County in the vicinity of Mist Gas Field, are the only federal lands presently under lease in Oregon. Total rental income to the BLM was \$11,760 during 1995. Applications were pending on 13,103 federal

acres at the end of 1995, which is a decrease from applications pending on 39,942 acres at the end of 1994 because the majority of such offers were withdrawn during the year.

The State of Oregon held no lease sales during 1995, and no State oil and gas leases were issued during the year. With no changes during the year, a total of 16 State of Oregon tracts comprising a total of 25,250 acres were under lease at year's end, and total rental income was \$25,250—the same as in 1994.

Columbia County held no lease sales during 1995.

DRILLING

For the second consecutive year, no exploratory oil and gas wells were drilled in Oregon. Before this, exploratory wells for oil and gas had been drilled in Oregon each year from 1974 until 1993. The lack of drilling during 1995 is largely attributed to the fact that Nahama and Weagant Energy Company, the operator of Mist Gas Field, had filed for bankruptcy during 1994. A court-appointed trustee was assigned to sell the assets of Nahama and Weagant Energy Company, including the Mist Gas Field. During 1995, Enerfin Resources of Houston, Texas, was the successful bidder for the Mist Gas Field and will become the operator of the field. It was reported that negotiations were completed during December 1995, which transferred ownership of ten wells and certain leases to Oregon Natural Gas Development Company (ONGD) of Portland, Oregon. ONGD is now the operator of the following wells: CC 13-34, CC 43-33-75, CC 23-22, CC 43-22, CC 31-27-65, CC 14-23, CC 44-27-65, CC 31-34-65, CC 21-35-65, and LF 12B-35-65. All other wells at Mist Gas Field are operated by Nahama and Weagant Energy pending finalization of the transfer of ownership to Enerfin Resources.

During 1995, DOGAMI did not issue any permits to drill, while four incomplete applications were withdrawn. Permit activity is listed in Table 1.

Table 1. Oil and gas permit activity in Oregon 1995

Permit number	Operator, well, API number	Location	Permit activity
440	Norwestco, Inc. 1-29 36-069-00009	SW¼ sec. 29 T. 9 S., R. 23 E. Wheeler County	Incomplete application, withdrawn.
485	Carbon Energy Menasha 28-1 36-001-00025	SW¼ sec. 28 T. 26 S., R. 13 W. Coos County	Incomplete application, withdrawn.
498	Carbon Energy Menasha 1-16 36-011-00028	SE¼ sec. 16 T. 26 S., R. 13 W. Coos County	Incomplete application, withdrawn.
499	Carbon Energy Menasha 1-16 36-011-00028	SE¼ sec. 32 T. 26 S., R. 13 W. Coos County	Incomplete application, withdrawn.

PRODUCTION

Despite the bankruptcy of Nahama and Weagant Energy Company, the Mist Gas Field operated normally during the year. The field had a total of 18 wells that were productive during the year, which is a decline from 21 wells productive during 1994. The wells produced 2.5 Bcf of gas, a significant decrease from the 4.2 Bcf of gas produced during 1994. This decrease was largely the result of the three factors. The first is the normal decline of the existing wells at the field. The second factor is the record low price received for gas at the Mist Gas Field since it was discovered during 1979, which discouraged exploration and production at the field. The gas prices paid during 1995 ranged from 4 cents to 11 cents per therm during the year until December, when they increased to 22 cents per therm for the month. During 1994, gas prices ranged from 11 to 23 cents per therm. The final reason for the production decline during 1995 was the decommissioning of the Nitrogen Rejection Unit (NRU) that was installed during 1994. The NRU enabled three low-Btu wells at Mist Gas Field to go on production by separating the methane gas from noncombustible nitrogen gas. However, due to the cost to run the NRU, decline in production of the low-Btu wells, and other factors, the NRU was decommissioned during the year and the three low-Btu wells were shut in. As a result of the low production volume and low gas prices during 1995, the total value of the gas produced at Mist Gas Field was \$1.8 million, a significant decline from the \$6.4 million during 1994. Cumulatively, the Mist Gas Field has produced about 56.5 Bcf of gas with a total value of \$112.8 million since it was discovered in 1979.

GAS STORAGE

The Mist Natural Gas Storage Project remained fully operational during 1995. The gas storage project has nine injection-withdrawal service wells, five in the Bruer Pool and four in the Flora Pool, and thirteen observation-monitor service wells. The two pools have a combined storage capacity of 10 Bcf of gas. This allows for the cycling of about 6 Bcf of gas in the reservoirs at pressures between approximately 400 and 1,000 psi and will provide for an annual delivery of 1 million therms of gas per day for 100 days. During 1995, about 4,469,288 cubic feet of gas was injected, and 3,566,865 cubic feet was withdrawn at the Mist Storage Project. Plans are underway by ONGD to develop additional storage pools at the Mist Gas Field during 1996.

OTHER ACTIVITIES

DOGAMI has completed a five-year study of the oil, gas, and coal resource potential of the Tye Basin located in Douglas and Coos Counties in the southern Coast Range. The study, which was funded by landowners in the study area and by county, state, and federal agencies in a public-private partnership, is an investigation of source rock, stratigraphy, and structural framework for those character-

(Continued on page 74)

ABSTRACTS OF PAPERS

The following abstract is of a paper given at an international conference in May 1995 at the University of Washington. The conference, titled "Tsunami deposits—geologic warnings of future inundation," was sponsored by the Quaternary Research Center, the National Oceanic and Atmospheric Administration, and the U.S. Geological Survey. Among the 80 registered participants were scientists from Canada, Germany, Japan, Norway, the Philippines, the United Kingdom, and the United States.

A paleohydraulic reconstruction of 300-year old tsunami deposits at Willapa Bay, Washington State, by Mary Ann Rinehart, GeoEngineers, Inc., Redmond, Wash.; and Joanne Bourgeois, Department of Geological Sciences, University of Washington

Sand sheets about 300 years old along the southern Washington coast can be explained by a tsunami generated at the Cascadia Subduction Zone but not by storm setup or seiche. These sand sheets, preserved in many parts of the interiors of Willapa Bay and Grays Harbor estuaries, overlie a buried peaty soil indicating abrupt subsidence. They are composed of one to five horizontal laminae of sandy silt or silty sand, alternating with laminae of mud. The character and sedimentology of the deposits indicate that (1) the depositional mechanism was a single, landward-directed event comprising multiple pulses; (2) each lamina was deposited rapidly from a high concentration of suspended sediment; (3) velocities and boundary shear stresses generated during each pulse were high enough to entrain and suspend sufficient volumes of sand and silt into the overbank part of the flow from tidal-channel floors; (4) overbank velocities were sufficient to transport sand up to about 200 m across marsh surfaces; and (5) a period of quiet water followed deposition of each sandy lamina.

As a test of the tsunami hypothesis we used a simple sediment-transport model to reconstruct the conditions necessary to deposit sandy laminae on marshes along the Niihau River, an arm of Willapa Bay. We assumed (conservatively) steady-flow conditions and uniform distribution of sand and silt throughout the overbank part of the water column. Discharge and suspended-sediment concentrations were reconstructed with a simple advection model applied to the lateral distribution of grain sizes in the basal sandy lamina. We then constrained the rate of lamina deposition, overbank flow depth, and pulse period by matching the component concentrations (estimated from the overbank trajectory model) to concentrations generated within the channel as calculated from the Rouse equation. The results of the reconstruction indicate overbank flow depths of 1–2 m, pulse periods of 18–30 min, and in-channel velocities of 340–400 cm/s.

A tsunami wave train provides the best fit to results of the reconstruction, based on wave period, number of pulses, water velocity, and distribution of the sand sheet around

Willapa Bay and Grays Harbor. The calculated wave periods are consistent with known tsunami periods and inconsistent with the calculated seiche period for Willapa Bay (1.5–2 hrs.)

The number of pulses is also improbable for a seiche, as it is unlikely the bay could seiche the required number of times, given its shallow, irregular bathymetry. Storms are unlikely to have produced the calculated flow velocities, maximum combined storm/tidal current velocities at the entrance of Willapa Bay are not known to exceed 250–300 cm/sec, and channel-flow velocities of that magnitude would not be maintained in protected tidal channels such as the Niawiakum River. Finally, the regional extent of the deposit, as well as its characteristics and local variation, can be explained by tsunami wave behavior. □

(Continued from page 73)

istics that are needed to generate and trap oil and gas. The final data, reports, and maps are expected to be published by DOGAMI during 1996. A series of maps and preliminary reports that present a revised understanding of the geologic framework of the Tyee Basin have already been published. Contact DOGAMI for a complete publication list including those for the Tyee Basin study.

The Northwest Energy Association, formerly the Northwest Petroleum Association, remained active for the year and has over 100 members. At its regular monthly meetings, speakers give talks generally related to energy matters in the Pacific Northwest. The 1995 symposium was held at Astoria, Oregon, on underground natural gas storage and energy-related matters of interest in the region. Plans are now underway for the 1996 symposium, which will be held in the Seattle, Washington, area. For information, contact the NWEA, P.O. Box 6679, Portland, OR 97228.

During 1996, no legislative or rule changes were made to DOGAMI statutes and administrative rules related to oil and gas drilling exploration activities. Copies of current statutes and administrative rules can be obtained from DOGAMI. □

In memoriam: Barbara Jacob, Cliff Speaker

The Oregon Department of Geology and Mineral Industries (DOGAMI) mourns the death of two former staff members, both retired.

Barbara Ann Jacob died March 4, 1996, at age 75, in Portland. She was a native of Tacoma and lived most of her life in Portland. She was a secretary for DOGAMI for 15 years until her retirement in 1986 and specialized in the regulatory work of the oil and gas section. Barbara leaves a son, R. Larry, in Vancouver, Washington; a daughter, Carolyn Harmon, in Olympia; a sister, Gloria Kingsley, in Milwaukie; a brother, Dee Swayze, in Santa Clara, California; and three grandchildren.

Clifford Drake Speaker died March 6, 1996, at age 82. He was born in Parkrose and lived in Vancouver, Washing-

DOGAMI PUBLICATIONS

Released May 1, 1996:

Tsunami Hazard Map of the Siletz Bay Area, Lincoln County, Oregon, by G.R. Priest, M. Qi, A.M. Baptista, C.D. Peterson, and M.E. Darienzo. Geological Map Series GMS-99, \$6.

This map of the area between Lincoln City and Glenden Beach outlines five tsunami hazard zones for elevations between sea level and about 100 ft (30 m). The map was produced by the joint efforts of scientists from DOGAMI, the Oregon Graduate Institute of Science and Technology, and Portland State University and is intended primarily for planning for evacuation in the event of a tsunami.

The tsunami hazard map shows a black-and-white air-photo image of the coastal zone with elevation contours superimposed. The scale of 1:12,000 (1 inch = 1,000 feet) allows identification of streets and medium-sized structures. Three different types of red lines mark tsunami runup elevations that serve to separate risk zones: extreme risk, high risk, moderate risk, and low risk; the fifth category, negligible risk, is identified as those lands above 100 feet (30 m) in elevation. The map also identifies drill sites where cores revealed buried soils and, in some cases, tsunami sand layers from prehistoric events.

For evacuation planning, the new tsunami map should be used in conjunction with the relative earthquake hazard maps of the area published earlier by DOGAMI as Geological Map Series map GMS-93 (\$20).

Explanation of Mapping Methods and Use of the Tsunami Hazard Map of the Siletz Bay Area, Lincoln County, Oregon, edited by G.R. Priest. DOGAMI Open-File Report O-95-05, 69 p., \$6.

This 69-page paper contains detailed discussions of the map mentioned above and of its use. It also describes the development of the methods used to produce the map and the geologic evidence of prehistoric earthquakes and tsunamis on the coast.

These DOGAMI publications are now available over the counter, by mail, FAX, or phone from the Nature of the Northwest Information Center and the DOGAMI field offices in Baker City and Grants Pass (addresses on p. 50). Orders may be charged to Visa or Mastercard. Orders under \$50 require prepayment. □

ton, from 1983 until his death. He was an accountant for Publishers Paper Company and then worked as business manager for DOGAMI from 1970 until his retirement in 1978. His surviving family includes his wife, Rachel C.; a daughter, Marylou S. Churchill, in Boston; sons Paul, in Upland, California, and Michael, in Portland; stepsons Roger Cole, in Vancouver, and Douglas Cole, in Bellingham, Washington; and three grandchildren. □

AVAILABLE PUBLICATIONS
OREGON DEPARTMENT OF GEOLOGY AND MINERAL INDUSTRIES

GEOLOGICAL MAP SERIES

Price ☑

GMS-5 Powers 15' quadrangle, Coos and Curry Counties. 1971	4.00
GMS-6 Part of Snake River canyon. 1974	8.00
GMS-8 Complete Bouguer gravity anomaly map, central Cascades. 1978	4.00
GMS-9 Total-field aeromagnetic anomaly map, central Cascades. 1978	4.00
GMS-10 Low- to intermediate-temperature thermal springs and wells. 1978	4.00
GMS-12 Oregon part, Mineral 15' quadrangle, Baker County. 1978	4.00
GMS-13 Huntington/Olds Ferry 15' quads., Baker/Malheur Counties. 1979	4.00
GMS-14 Index to published geologic mapping in Oregon, 1898-1979. 1981	8.00
GMS-15 Gravity anomaly maps, north Cascades. 1981	4.00
GMS-16 Gravity anomaly maps, south Cascades. 1981	4.00
GMS-17 Total-field aeromagnetic anomaly map, south Cascades. 1981	4.00
GMS-18 Rickreall, Salem West, Monmouth, and Sidney 7½' quadrangles, Marion and Polk Counties. 1981	6.00
GMS-19 Bourne 7½' quadrangle, Baker County. 1982	6.00
GMS-20 S½ Burns 15' quadrangle, Harney County. 1982	6.00
GMS-21 Vale East 7½' quadrangle, Malheur County. 1982	6.00
GMS-22 Mount Ireland 7½' quadrangle, Baker/Grant Counties. 1982	6.00
GMS-23 Sheridan 7½' quadrangle, Polk and Yamhill Counties. 1982	6.00
GMS-24 Grand Ronde 7½' quadrangle, Polk/Yamhill Counties. 1982	6.00
GMS-25 Granite 7½' quadrangle, Grant County. 1982	6.00
GMS-26 Residual gravity, north/central/south Cascades. 1982	6.00
GMS-27 Geologic and neotectonic evaluation of north-central Oregon. The Dalles 1° × 2° quadrangle. 1982	7.00
GMS-28 Greenhorn 7½' quadrangle, Baker/Grant Counties. 1983	6.00
GMS-29 NE¼ Bates 15' quadrangle, Baker/Grant Counties. 1983	6.00
GMS-30 SE¼ Pearsoll Peak 15' quad., Curry/Josephine Counties. 1984	7.00
GMS-31 NW¼ Bates 15' quadrangle, Grant County. 1984	6.00
GMS-32 Wilhoit 7½' quadrangle, Clackamas/Marion Counties. 1984	5.00
GMS-33 Scotts Mills 7½' quad., Clackamas/Marion Counties. 1984	5.00
GMS-34 Stayton NE 7½' quadrangle, Marion County. 1984	5.00
GMS-35 SW¼ Bates 15' quadrangle, Grant County. 1984	6.00
GMS-36 Mineral resources of Oregon. 1984	9.00
GMS-37 Mineral resources, offshore Oregon. 1985	7.00
GMS-38 NW¼ Cave Junction 15' quadrangle, Josephine County. 1986	7.00
GMS-39 Bibliography and index: ocean floor, continental margin. 1986	6.00
GMS-40 Total-field aeromagnetic anomaly maps, northern Cascades. 1985	5.00
GMS-41 Elkhorn Peak 7½' quadrangle, Baker County. 1987	7.00
GMS-42 Ocean floor off Oregon and adjacent continental margin. 1986	9.00
GMS-43 Eagle Butte & Gateway 7½' quads., Jefferson/Wasco C. 1987	5.00
as set with GMS-44 and GMS-45	11.00
GMS-44 Seekseequa Junct./Metolius B. 7½' quads., Jefferson C. 1987	5.00
as set with GMS-43 and GMS-45	11.00
GMS-45 Madras West/East 7½' quads., Jefferson County. 1987	5.00
as set with GMS-43 and GMS-44	11.00
GMS-46 Breitenbush River area, Linn and Marion Counties. 1987	7.00
GMS-47 Crescent Mountain area, Linn County. 1987	7.00
GMS-48 McKenzie Bridge 15' quadrangle, Lane County. 1988	9.00
GMS-49 Map of Oregon seismicity, 1841-1986. 1987	4.00
GMS-50 Drake Crossing 7½' quadrangle, Marion County. 1986	5.00
GMS-51 Elk Prairie 7½' quadrangle, Marion and Clackamas Counties. 1986	5.00
GMS-52 Shady Cove 7½' quadrangle, Jackson County. 1992	6.00
GMS-53 Owyhee Ridge 7½' quadrangle, Malheur County. 1988	5.00
GMS-54 Graveyard Point 7½' quad., Malheur/Owyhee Counties. 1988	5.00
GMS-55 Owyhee Dam 7½' quadrangle, Malheur County. 1989	5.00
GMS-56 Adrian 7½' quadrangle, Malheur County. 1989	5.00
GMS-57 Grassy Mountain 7½' quadrangle, Malheur County. 1989	5.00
GMS-58 Double Mountain 7½' quadrangle, Malheur County. 1989	5.00
GMS-59 Lake Oswego 7½' quad., Clackam., Multn., Wash. Counties. 1989	7.00
GMS-60* Damascus 7½' quad., Clackam., Multn. Counties. 1994	8.00
GMS-61 Mitchell Butte 7½' quadrangle, Malheur County. 1990	5.00
GMS-62* The Elbow 7½' quadrangle, Malheur County. 1993	8.00
GMS-63 Vines Hill 7½' quadrangle, Malheur County. 1991	5.00
GMS-64 Sheaville 7½' quadrangle, Malheur County. 1990	5.00
GMS-65 Mahogany Gap 7½' quadrangle, Malheur County. 1990	5.00
GMS-66 Jonesboro 7½' quadrangle, Malheur County. 1992	6.00
GMS-67 South Mountain 7½' quadrangle, Malheur County. 1990	6.00
GMS-68 Reston 7½' quadrangle, Douglas County. 1990	6.00
GMS-69 Harper 7½' quadrangle, Malheur County. 1992	5.00
GMS-70 Boswell Mountain 7½' quadrangle, Jackson County. 1992	7.00
GMS-71 Westfall 7½' quadrangle, Malheur County. 1992	5.00
GMS-72 Little Valley 7½' quadrangle, Malheur County. 1992	5.00

Price ☑

GMS-73* Cleveland Ridge 7½' quadrangle, Jackson County. 1993	5.00
GMS-74 Namorf 7½' quadrangle, Malheur County. 1992	5.00
GMS-75 Portland 7½' quadrangle, Multn., Wash., Clark Counties. 1991	7.00
GMS-76 Carnas Valley 7½' quadrangle, Douglas and Coos Counties. 1993	6.00
GMS-77 Vale 30×60 minute quadrangle, Malheur County. 1993	10.00
GMS-78 Mahogany Mountain 30×60 minute quadrangle, Malheur C. 1993	10.00
GMS-79* Earthquake hazards, Portland 7½' quad., Multnomah C. 1993	20.00
GMS-80* McLeod 7½' quadrangle, Jackson County. 1993	5.00
GMS-81* Tumalo Dam 7½' quadrangle, Deschutes County. 1994	6.00
GMS-82* Limber Jim Creek 7½' quadrangle, Union County. 1994	5.00
GMS-83* Kenyon Mountain 7½' quadrangle, Douglas/Coos Counties. 1994	6.00
GMS-84* Remote 7½' quadrangle, Coos County. 1994	6.00
GMS-85* Mount Gurney 7½' quadrangle, Douglas/Coos Counties. 1994	6.00
GMS-86* Tenmile 7½' quadrangle, Douglas County. 1994	6.00
GMS-87* Three Creek Butte 7½' quadrangle, Deschutes County. 1995	6.00
GMS-88* Lakecreek 7½' quadrangle, Jackson County. 1995	8.00
GMS-89* Earthquake hazards, Mt. Tabor 7½' quad., Multnomah C. 1995	10.00
GMS-90* Earthquake hazards, Beaverton 7½' quad., 1995	10.00
GMS-91* Earthquake hazards, Lake Oswego 7½' quad., 1995	10.00
GMS-92* Earthquake hazards, Gladstone 7½' quad., 1995	10.00
GMS-93* Earthquake hazards, Siletz Bay area, Lincoln County, 1995	20.00
GMS-94* Charleston 7½' quadrangle, Coos County. 1995	8.00
GMS-97* Coos Bay 7½' quadrangle, Coos County. 1995	6.00
GMS-98* Dora and Sitkum 7½' quadrangles, Coos County. 1995	6.00
GMS-99* Tsunami hazard map, Siletz Bay area, Lincoln County. 1995	6.00

SPECIAL PAPERS

2 Field geology, SW Broken Top quadrangle. 1978	5.00
3 Rock material resources, Clackam., Columb., Multn., Wash. C. 1978	8.00
4 Heat flow of Oregon. 1978	4.00
5 Analysis and forecasts of demand for rock materials. 1979	4.00
6 Geology of the La Grande area. 1980	6.00
7 Pluvial Fort Rock Lake, Lake County. 1979	5.00
8 Geology and geochemistry of the Mount Hood volcano. 1980	4.00
9 Geology of the Breitenbush Hot Springs quadrangle. 1980	5.00
10 Tectonic rotation of the Oregon Western Cascades. 1980	4.00
11 Bibliography and index of theses and dissertations, 1899-1982. 1982	7.00
12 Geologic linears, northern part of Cascade Range, Oregon. 1980	4.00
13 Faults and lineaments of southern Cascades, Oregon. 1981	5.00
14 Geology and geothermal resources, Mount Hood area. 1982	8.00
15 Geology and geothermal resources, central Cascades. 1983	13.00
16 Index to <i>Ore Bin</i> (1939-78) and <i>Oregon Geology</i> (1979-82). 1983	5.00
17 Bibliography of Oregon paleontology, 1792-1983. 1984	7.00
18 Investigations of talc in Oregon. 1988	8.00
19 Limestone deposits in Oregon. 1989	9.00
20 Bentonite in Oregon. 1989	7.00
21 Field geology, NW¼ Broken Top 15' quadrangle, Deschutes C. 1987	6.00
22 Silica in Oregon. 1990	8.00
23 Forum on Geology of Industrial Minerals, 25th, 1989, Proceedings. 1990	10.00
24 Index to Forums on the Geology of Industrial Minerals, 1965-1989. 1990	7.00
25 Pumice in Oregon. 1992	9.00
26 Onshore-offshore geol. cross section, N. Coast Range to cont. slope. 1992	11.00
27 Economic analysis, construction aggregate markets and forecast. 1995	15.00

OIL AND GAS INVESTIGATIONS

3 Foraminifera, General Petroleum Long Bell #1 well. 1973	4.00
Foraminifera, E.M. Warren Coos County 1-7 well. 1973	4.00
5 Prospects for natural gas, upper Nehalem River Basin. 1976	6.00
6 Prospects for oil and gas, Coos Basin. 1980	10.00
7 Correlation of Cenozoic stratigraphic units, W. Oregon/Washington. 1983	9.00
8 Subsurface stratigraphy of the Ochoco Basin, Oregon. 1984	8.00
9 Subsurface biostratigraphy of the east Nehalem Basin. 1983	7.00
10 Mist Gas Field: Exploration/development, 1979-1984. 1985	5.00
11 Biostratigraphy of exploratory wells, W. Coos, Douglas, Lane Co. 1984	7.00
12 Biostratigraphy, exploratory wells, N. Willamette Basin. 1984	7.00
13 Biostratigraphy, exploratory wells, S. Willamette Basin. 1985	7.00
14 Oil and gas investigation of the Astoria Basin. 1985	8.00
15 Hydrocarbon exploration and occurrences in Oregon. 1989	8.00
16 Available well records and samples, onshore/offshore. 1987	6.00
17 Onshore-offshore cross section, Mist Gas Field to cont. shelf/slope. 1990	10.00
18 Schematic fence diagram, S. Tyee basin, Oregon Coast Range. 1993	9.00

AVAILABLE DEPARTMENT PUBLICATIONS (continued)

BULLETINS

	Price <input checked="" type="checkbox"/>
33 Bibliography, geol. & min. res. of Oregon (1st suppl. 1936-45). 1947	4.00
36 Papers on Tertiary Foraminifera (v. 2 [parts VII-VIII] only). 1949	4.00
44 Bibliography (2nd supplement, 1946-50). 1953	4.00
46 Ferruginous bauxite, Salem Hills, Marion County. 1956	4.00
53 Bibliography (3rd supplement, 1951-55). 1962	4.00
65 Proceedings of the Andesite Conference. 1969	11.00
67 Bibliography (4th supplement, 1956-60). 1970	4.00
71 Geology of lava tubes, Bend area, Deschutes County. 1971	6.00
78 Bibliography (5th supplement, 1961-70). 1973	4.00
82 Geologic hazards of Bull Run Watershed, Multn./Clackam. C. 1974	8.00
87 Environmental geology, western Coos/Douglas Counties. 1975	10.00
88 Geology/min. res., upper Chetco R. drainage, Curry/Josephine C. 1975	5.00
89 Geology and mineral resources of Deschutes County. 1976	8.00
90 Land use geology of western Curry County. 1976	10.00
91 Geologic hazards, parts of N. Hood River, Wasco, Sherman C. 1977	9.00
92 Fossils in Oregon. Collection of reprints from the <i>Ore Bin</i> . 1977	5.00
93 Geology, mineral resources, and rock material, Curry County. 1977	8.00
94 Land use geology, central Jackson County. 1977	10.00
95 North American ophiolites (IGCPproject). 1977	8.00
96 Magma genesis. AGU Chapman Conf. on Partial Melting. 1977	15.00
97 Bibliography (6th supplement, 1971-75). 1978	4.00
98 Geologic hazards, eastern Benton County. 1979	10.00
99 Geologic hazards of northwestern Clackamas County. 1979	11.00
101 Geologic field trips in W. Oregon and SW. Washington. 1980	10.00
102 Bibliography (7th supplement, 1976-79). 1981	5.00
103 Bibliography (8th supplement, 1980-84). 1987	8.00

MISCELLANEOUS PAPERS

5 Oregon's gold placers. 1954	2.00
11 Articles on meteorites (reprints from the <i>Ore Bin</i>). 1968	4.00
15 Quicksilver deposits in Oregon. 1971	4.00
19 Geothermal exploration studies in Oregon, 1976. 1977	4.00
20 Investigations of nickel in Oregon. 1978	6.00

SHORT PAPERS

25 Petrography of Rattlesnake Formation at type area. 1976	4.00
27 Rock material resources of Benton County. 1978	5.00

MISCELLANEOUS PUBLICATIONS

Relative earthquake hazard map, Portland quadrangle (DOGAMI/Metro), 1993, with scenario report (add \$3.00 for mailing)	10.00
Geology of Oregon, 4th ed., E.L. and W.N. Orr and E.M. Baldwin, 1991, published by Kendall/Hunt (add \$3.00 for mailing)	26.95
Geologic map of Oregon, G.W. Walker and N.S. MacLeod, 1991, published by USGS (add \$3.00 for mailing)	11.50
Geological highway map, Pacific Northwest region, Oregon, Washington, and part of Idaho (published by AAPG). 1973	8.00
Oregon Landsat mosaic map (published by ERSAL, OSU). 1983	11.00
Mist Gas Field map, rev. 1995, with 1993-94 production figs. (OFR O-95-1)	8.00
Digital form of map (CAD formats .DGN, .DWG, .DXF), 3½-in. diskette	25.00
Mist Gas Field production figures 1979 through 1992 (OFR O-94-6)	5.00
Northwest Oregon, Correlation Sec. 24. Bruer & others, 1984 (AAPG)	6.00
Oregon rocks and minerals, a description. 1988 (OFR O-88-6)	6.00
Mineral information layer for Oregon by county (MILOC), 1993 update (OFR O-93-8), 2 diskettes (5¼-in., high-density, MS-DOS)	25.00
Directory of mineral producers, 1993 update, 56 p. (OFR O-93-9)	8.00
Geothermal resources of Oregon (published by NOAA). 1982	4.00
Mining claims (State laws governing quartz and placer claims)	Free
Back issues of <i>Oregon Geology</i>	3.00
Color postcard with Oregon State Rock and State Gemstone	1.00

Separate price lists for open-file reports, tour guides, recreational gold mining information, and non-Departmental maps and reports will be mailed upon request. **GMS maps marked with an asterisk (*) are available in digital form on diskette (geological information only).**

The Department also sells Oregon topographic maps published by the U.S. Geological Survey.

ORDER AND RENEWAL FORM

Check desired publications in list above or indicate how many copies and enter total amount below. Send order to **The Nature of the Northwest Information Center, Suite 177, 800 NE Oregon Street, Portland, OR 97232-2162**, or FAX (503) 731-4066. If you wish to order by phone, have your credit card ready and call (503) 872-2750. Payment must accompany orders of less than \$50. Payment in U.S. dollars only. Publications are sent postpaid. All sales are final. Subscription price for *Oregon Geology*: \$10 for 1 year, \$22 for 3 years.

Renewal ___ / new subscription ___ to *Oregon Geology*: 1 year (\$10) or 3 years (\$22) \$ _____
 Total amount for publications marked above: \$ _____
 Total payment enclosed or to be charged to credit card as indicated below: \$ _____

Name _____
 Address _____
 City/State/Zip _____

Please charge to Visa ___ / Mastercard ___, account number: _____

Expiration date: _____

Cardholder's signature _____